# HSOA Journal of Angiology & Vascular Surgery

## Validation of Caprini's Saphenous Ablation Scores

## J Caprini<sup>1</sup>, Fakhry A<sup>2\*</sup> and Nagib S<sup>3</sup>

<sup>1</sup>Pritzker School of Medicine, North Shore University Health System Evanston, USA

<sup>2</sup>Department of Vascular Surgery, Egyptian Military Academy, Egypt

<sup>3</sup>Department of Vascular Surgery, Alexandria Armed Forces Hospital, Egypt

## Abstract

**Introduction:** Deep Vein Thrombosis (DVT) after varicose vein surgery is well recognized. Less well documented is Endovenous Heat-Induced Thrombosis (EHIT), thrombus extension into a deep vein after superficial venous thermo ablation. There is no current agreement on the routine use of thromboprophylaxis in patients undergoing varicose vein surgery. more data on the incidence of VTE, and the need for postoperative thromboprophylaxis are necessary to formulate evidence-based clinical guidelines.

**Aim of the study:** Comparison of the use of Caprini's Saphenous Ablation Scores versus Standard Caprini's Score for the prophylaxis of VTE in EVLA.

**Patients & Methods:** 60 patients admitted to Royal Vascular Center, Alexandria - Egypt. From Jan. 1st, 2021, to June 30th 2021 divided in 3 groups. - Group A) Treated by ablation of Great saphenous vein and received VTE prophylaxis according to Caprini's Saphenous Ablation Scores Fig 1-A & B -Group B) Treated by ablation of Great saphenous vein and received VTE prophylaxis according to Standard Caprini's Risk Assessment Score for VTE Fig 2-A&B - C) Treated by ablation of Great saphenous vein without VTE prophylaxis Ablation was done using radial 1480 YAG laser fibers and Post-operative Duplex was done one day, 3 month and one-year.

**Results:** Age: 35.6+0.2, 34.3+0.5 and 37.6+0.09 years in the 3 groups M/F: 6/14, 8/12 and 7/13. In group A) all patients showed absent reflux post operatively and in all follow up visits. Duplex examination revealed significant reflux in the great saphenous veins in all patients and ranged from 0.7:0.9 Sec. and mean duration 0.7+0.09 Sec. Vein diameter ranged from (0.6: 0.9,X+0.76 CM.) in group A), (0.6: 0.8,X+0.7 CM.) in group B), (0.5: 0.8,X+0.65 CM.) in

\*Corresponding author: Fakhry A, Department of Vascular Surgery, Egyptian Military Academy, Egypt; Tel: +20 1223779844; E-mail: ayman\_vasc@live.com

**Citation:** Caprini J, Fakhry A, Nagib S (2021) Validation of Caprini's Saphenous Ablation Scores. J Angiol Vasc Surg 6: 082.

Received: December 20, 2021; Accepted: December 21, 2021; Published: December 28, 2021

**Copyright:** © 2021 J Caprini, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

group C) According to Caprini's saphenous score 14 patients 70% were classified as low risk, 4 patients 20% were of moderate risk and 2 patients 10% of high-risk in Group A, while applying standard Caprini's Score in group B patients revealed: 6 patients 30% were of moderate risk and 14 patients 70% of high risk in Group B). 3 patients developed superficial thrombophlebitis 15% and one patient 5% developed left calf DVT in Group C, while no VTE events were shown in both A&B Groups Significant drop in CVSS in patients in the three groups.

**Conclusion:** Risk assessment of patients undergoing varicose veins ablation should be done and applying Caprini's saphenous score as a better alternative to Standard Caprini's score for VTE prophylaxis and is considered a safe and cost-effective tool in VTE prevention in these patients.

**Keywords:** Risk assessment score; Thrombosis prophylaxis; Varicose veins; Venous thromboembolism

## Introduction

Laser Ablation of Varicose vein is among the most performed procedures worldwide, Venous Thromboembolism (VTE), either as Deep Venous Thrombosis (DVT) or a potentially fatal Pulmonary Embolus (PE), is the most serious complication in all of these procedures after treatment of varicose veins [1-3]. Hilly, et al. pointed out that there is no consensus of the perioperative thrombosis prophylaxis after varicose veins laser ablation [4]. Clinical guidelines by the American ACCP, Scottish SIGN and Dutch CBO all recommend risk stratification after which only patients with risk factors for thrombosis are to be treated with prophylaxis. The most important risk factors are obesity and a history of VTE [5].

## Aim of the Study

Comparison of the use of Caprini's Saphenous Ablation Scores versus Standard Caprini's Score for the prophylaxis of VTE in EVLA.

## **Patients & Methods**

60 patients admitted to Royal Vascular Center, Alexandria - Egypt. From Jan. 1st 2021 to June 30th 2021 divided in 3 groups. - Group A) Treated by ablation of Great saphenous vein and received VTE prophylaxis according to Caprini's Saphenous Ablation Scores Figure 1 A &B -Group B) Treated by ablation of Great saphenous vein and received VTE prophylaxis according to Standard Caprini's Risk Assessment Score for VTE Figure 2 C) Treated by ablation of Great saphenous vein without VTE prophylaxis Patients signed an informed consent [6,7]. The pre-operative assessment by venous Duplex Ultrasound to assess the incompetence of SFJ and DVT detection prior to intervention. Ablation was done using radial 1470 YAG laser fibers and Post-operative Duplex was done one day and one month.

## **Inclusion Criteria**

Patients with primary varicose veins with saphenofemoral junction incompetence with CEAP classification 2-6.



Figure 1: A & B) Caprini's Saphenous Ablation.



		The second second state from the second second from the Second state second		
۰	Very low risk (+0.5%)	No specific pharmacologic (unade Hit) or mechanical (unade 24.) prophysiki de used other than early ambulation		
1-2	Low risk (-1.5%)	Mechanical prophylaxis, preferably with PC (Grade 2C)		
ы	Moderalie risk (-7%)	(JMMH (Srade 28), LDUH (Srade 28) or mechanical prophylais with IPC (Srade 2C)	Mechanical prophylaxis, preferably with PC (Grade 2C)	
đ	High risk (-d1)	(3M8H (Srade 18) or LDUH (Srade 18) plus mechanical prophytiks with ES or IPC (Srade 2C)	Mechanical prophytals, preferably with PC_ with the test of baseding derivatives and phomocologic prophytales may be informed (Prode 2C)	
	High risk Concer surgery (stodominal or petivic)	UMIH (Srade 18) or LDUH (Srade 18) plus mechanical prophytois with (S or IPC (Srade 2C) Entended-duration prophytois (il weeks) with LMIM portdischarge (Grade 18)		
	High risk UMM-and LDUH contraindicated	Fondaparinus (Grade 2C) or low-dose aspirin (Febmg) (Grade 2C); mechanical (prophyliaik, preferably		

Figure 2: Standard Caprini's Risk Assessment.

## **Exclusion Criteria**

- Patients with history of DVT
- · Patients with recurrent varicose veins
- · Patients who are unable to provide an informed consent
- · Pregnant patients at the time of the examination

#### Results

Age: 35.6+0.2, 34.3+0.5 and 37.6+0.09 years in the 3 groups M/F: 6/14, 8/12 and 7/13. In group A) all patients showed absent reflux post operatively and in all follow up visits .Dilated veins was noticed on the left side in 12 patients 60%, 2 patient on right and on 6 patients 30 % in Group (A), 14 patients 70%, one patient 5% and 5 patients 25% in Group (B) and 13 patients 65%, 2 patients 10% and 5 patients 25% in Group (C), Table 1.

	Left side	Right side	Both sides	
Group A	12 patients 60%	2 patients 10%	6 patients 30%	
Group B	14 patients 70%	one patient 5%	5 patients 25%	
Group C	13 patients 65%	2 patients 10%	5 patients 25%	
Table 1: Side of varicose veins in the groups.				

Clinical presentation according to CEAP Classification showed that: (2,8,9,0,1 patents) were C2,3,4,5,and 6 in Group A , (2,7,9,1,1 patents) were C2,3,4,5,and 6 in Group B and (3,6,10,0,1 patents) were C2,3,4,5,and 6 in Group C. Duplex examination revealed significant reflux in the great saphenous veins in all patients and ranged from 0.7:0.9 Sec. and mean duration 0.7+0.09 Sec. Vein diameter ranged from (0.6: 0.9,X+0.76 CM.) in group A), (0.6: 0.8,X+0.7 CM.) in group B) , (0.5: 0.8,X+0.65 CM.) in group C) Table 2.

	Group A	Group B	Group C		
GSV diameter	Range:0.7:0.9 CM X 0.7+0.09 CM	Range:0.7:0.9 CM X 0.7+0.09 CM	Range:0.7:0.9 CM X 0.7+0.09 CM		
Table 2: Preoperative Duplex.					

According to Caprini's saphenous score 14 patients 70% were classified as low risk who used compression stockings only , 4 patients 20% were of moderate risk who received 40mg Enoxaparin / day for one week in addition to the stockings and 2 patients 10% of high-risk who received 40mg Enoxaparin /day for 28 days in addition to the stockings in Group A, while applying standard Caprini's Score in group B patients revealed : 6 patients 30% were of moderate risk and 14 patients 70% of high risk in Group B ) Table 3.

	Group A	Group B
Low risk	14 patients 70%	ZERO
Moderate risk	4 patients 20%	6 patients 30%
High risk	2 patients 10%	14 patients 70%

 Table 3: Risk assessment in Group A &B patients according to both risk assessment scores.

So, 14 patients 70% received Enoxaparin 40 mg / day for 7 days in group B patients, only 4 patients 20% received the same dose and duration ,with 2 patients 10% received the same dose for 28 days in group A patients. No VTE events were shown in both A&B, while 3 patients developed superficial thrombophlebitis 15% and one patient Citation: Caprini J, Fakhry A, Nagib S (2021) Validation of Caprini's Saphenous Ablation Scores. J Angiol Vasc Surg 6: 082.

5% developed left calf DVT in Group C, and the estimated Saphenous score was 11 and the standard score was 7 for these patients. Significant drop in VCSS in patients in the three groups, it dropped from 10.9 to 8.1, from 10.7 to 7.9 and 10.6 to 7.7 in group A, B and C. The time needed to complete total ablation of GSV ranged from 40:51.5, with a mean time 43.6+0.3, 44.3+0.9, 43.7+0.75 minutes the 3 groups.

So, 14 patients 70% received Enoxaparin 40 mg / day for 7 days in group B patients, only 4 patients 20% received the same dose and duration ,with 2 patients 10% received the same dose for 28 days in group A patients. No VTE events were shown in both A&B, while 3 patients developed superficial thrombophlebitis 15% and one patient 5% developed left calf DVT in Group C, and the estimated Saphenous score was 11 and the standard score was 7 for these patients. Significant drop in VCSS in patients in the three groups, it dropped from 10.9 to 8.1, from 10.7 to 7.9 and 10.6 to 7.7 in group A, B and C. The time needed to complete total ablation of GSV ranged from 40:51.5, with a mean time 43.6+0.3, 44.3+0.9, 43.7 + 0.75 minutes the 3 groups [8-10].

## Conclusion

Risk assessment of patients undergoing varicose veins ablation should be done and applying Caprini's saphenous score as a better alternative to Standard Caprini's score for VTE prophylaxis and is considered a safe and cost-effective tool in VTE prevention in these patients.

## Disclosure

The authors have no conflict of interest to declare.

## **Conflicts of Interest**

The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

## Funding

The authors report no involvement in the research by the sponsor that could had influenced the outcome of this work.

## Acknowledgements

The authors acknowledge doctors and nursing staff in vascular surgery department of Alexandria Armed Forces hospital for proofreading activity.

## References

- Marsh P, Price BA, Holdstock J, Harrison C, Whiteley MS (2010) Deep Vein Thrombosis (DVT) after Venous Thermoablation Techniques: Rates of Endovenous Heat-induced Thrombosis (EHIT) and Classical DVT after Radiofrequency and Endovenous Laser Ablation in a Single Centre. Eur J Vasc Endovasc Surg 40: 521-272.
- Testroote MJG, Wittens CHA (2013) Prevention of venous thromboenbolism in patients undergoing surgical treatment of varicose veins. Phlebology 1: 86-90.
- Dexter D, Kabnick L, Berland T, Jacobowitz G, Lamparello P, et al. (2012) Complications of endovenous lasers. Phlebology 27: 40-45.
- 4. Healy DA, Kimura S, Power D, Alhaj A, Abdeldaim Y, et al. (2018) A Systematic Review and Meta-analysis of Thrombotic Events Following Endovenous Thermal Ablation of the Great Saphenous Vein. Eur J Vasc Endovasc Surg 56: 410-424.
- Jacobs CE, Pinzon MM, Orozco J, Hunt PJB, Rivera A, et al. (2014) Deep venous thrombosis after saphenous endovenous radiofrequency ablation: Is it predictable? Ann Vasc Surg 28: 679-685.
- Caprini JA (2021) Thrombosis Prophylaxis in Varicose Vein Patients, Patient Evaluation and Degree of Risk. Presented online. Egyptian Venous Forum.
- Cronin MA, Dengler N, Krauss ES, Segal A, Wei N, et al. (2019) Completion of the Updated Caprini Risk Assessment Model (2013 Version). Clin Appl Thromb Hemost 25: 1-10.
- Mao J, Zhang C, Wang Z, Gan S, Ll K (2012) A retrospective study comparing endovenous laser ablation and microwave ablation for great saphenous varicose veins. Eur Rev Med Pharmacol Sci 16: 873-877.
- 9. Shutze WP, Kane K, Hicks T, Kedora J, Hohmann S, et al. (2021) Bertram Smith Baylor University Medical Center, Dallas, TX . The Incidence and Outcome of DVT after Endovenous Laser Ablation.
- Enoch S, Woon E, Blair SD (2003) Thromboprophylaxis can be omitted in selected patients undergoing varicose vein surgery and hernia repair. Br J Surg 90: 818-820.

Page 3 of 3



Advances In Industrial Biotechnology | ISSN: 2639-5665 Advances In Microbiology Research | ISSN: 2689-694X Archives Of Surgery And Surgical Education | ISSN: 2689-3126 Archives Of Urology Archives Of Zoological Studies | ISSN: 2640-7779 Current Trends Medical And Biological Engineering International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276 Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292 Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370 Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594 Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562 Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608 Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879 Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397 Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751 Journal Of Aquaculture & Fisheries | ISSN: 2576-5523 Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780 Journal Of Biotech Research & Biochemistry Journal Of Brain & Neuroscience Research Journal Of Cancer Biology & Treatment | ISSN: 2470-7546 Journal Of Cardiology Study & Research | ISSN: 2640-768X Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943 Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771 Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844 Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801 Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978 Journal Of Cytology & Tissue Biology | ISSN: 2378-9107 Journal Of Dairy Research & Technology | ISSN: 2688-9315 Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783 Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798 Journal Of Environmental Science Current Research | ISSN: 2643-5020 Journal Of Food Science & Nutrition | ISSN: 2470-1076 Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566

Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485 Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662 Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999 Journal Of Hospice & Palliative Medical Care Journal Of Human Endocrinology | ISSN: 2572-9640 Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654 Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493 Journal Of Light & Laser Current Trends Journal Of Medicine Study & Research | ISSN: 2639-5657 Journal Of Modern Chemical Sciences Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044 Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313 Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400 Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419 Journal Of Obesity & Weight Loss | ISSN: 2473-7372 Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887 Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052 Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X Journal Of Pathology Clinical & Medical Research Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649 Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670 Journal Of Plant Science Current Research | ISSN: 2639-3743 Journal Of Practical & Professional Nursing | ISSN: 2639-5681 Journal Of Protein Research & Bioinformatics Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150 Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177 Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574 Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060 Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284 Journal Of Toxicology Current Research | ISSN: 2639-3735 Journal Of Translational Science And Research Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193 Journal Of Virology & Antivirals Sports Medicine And Injury Care Journal | ISSN: 2689-8829 Trends In Anatomy & Physiology | ISSN: 2640-7752

## Submit Your Manuscript: https://www.heraldopenaccess.us/submit-manuscript