

## Research Article

# Seasonal Variations in Hematological and Serum Biochemical Parameters in Kiko Meat Goats under Semi-Intensive Management Systems

Cokere Okere<sup>1\*</sup>, King R<sup>1</sup> and Gurung N<sup>2</sup>

<sup>1</sup>Department of Agricultural and Environmental Sciences, Tuskegee University, Tuskegee, Alabama, USA

<sup>2</sup>George Washington Carver Agricultural Experiment Station, Tuskegee University, Tuskegee, Alabama, USA

### Abstract

Hematological and serum biochemical profiles have been used in several species of domestic livestock to monitor herd health. Application of this technique to meat goats in semi-intensive production systems has been limited by a lack of suitable reference ranges for most of these parameters. Mature Kiko meat goats (n = 15; age = 9-12 months) were utilized to determine the effects of season on biochemical parameters and hematology profiles. Meteorological data (rainfall, diurnal temperature and relative humidity) were also collected. For hematological profiles, non-significant seasonal differences were observed in hemoglobin (HGB) values of  $9.67 \pm 0.75$ ,  $9.23 \pm 1.478$ ,  $8.2 \pm 1.21$  and  $9.99 \pm 1.59$  G/DL  $P \geq 0.05$  for spring, summer, fall and winter respectively. Red Blood Cells (RBC) were significantly affected ( $P \leq 0.05$ ) by season of collection  $16.7 \pm 4.12$ ,  $18.1 \pm 2.52$ ,  $18.0 \pm 1.58$  and  $20.5 \pm 4.27$  M/ $\mu$ L for spring, summer, fall and winter respectively. Also, White Blood Cells (WBC) showed non-significant seasonal variations ( $P \geq 0.05$ )  $15.04 \pm 3.53$ ,  $15.9 \pm 6.47$ ,  $14.3 \pm 5.02$ , and  $14.8 \pm 6.59$  K/ $\mu$ L for spring, summer, fall and winter respectively. However, highly significant ( $P \leq 0.01$ ) seasonal differences were observed Reticulocytes (RETIC) K/ $\mu$ L  $3.76 \pm 2.29$ ,  $1.34 \pm 2.0$ ,  $0.90 \pm 1.26$  and  $1.56 \pm 1.49$  for spring, summer, fall and winter respectively. Serum biochemical parameters showed non-significant blood glucose values of  $65.0 \pm 6.2$ ,  $62.7 \pm 6.2$ ,  $60.0 \pm 16.6$  and  $57.8 \pm 11.8$  mg/dL  $P \geq 0.05$  for spring, summer, fall and winter respectively.

\*Corresponding author: Cokere Okere, Department of Agricultural and Environmental Sciences, Tuskegee University, Tuskegee, Alabama, E-mail: cokere@tuskegee.edu

**Citation:** Okere C, King R, Gurung N (2022) Seasonal Variations in Hematological and Serum Biochemical Parameters in Kiko Meat Goats under Semi-Intensive Management Systems. J Anim Res Vet Sci 6: 0.37.

**Received:** September 17, 2022; **Accepted:** September 30, 2022; **Published:** October 7, 2022

**Copyright:** © 2022. Okere C, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Blood urea nitrogen (BUN mg/DL), Calcium (CA mg/DL) and Alkaline Aminotransferase ALT U/L showed a high significant seasonal variation ( $P \leq 0.01$ ). The hematological and serum biochemical values obtained in this study will further underlined the need to establish appropriate physiological baseline values for meat goats in Alabama which could help in realistic evaluation of the management practice, nutrition, diagnosis of health as well as in determining the physiological status of goats among limited resource producers.

**Keywords:** Blood serum biochemistry; Hematology; Kiko meat goats; Seasons

### Abbreviations

BWT: Body Weight

BCS: Body Condition Score

GLU: Glucose

CREA: Creatinine

BUN: Blood Urea Nitrogen

CA: Calcium

TP: Total Protein

ALT: Alkaline Aminotransferase

AST: Aspartate Aminotransferase

ALKP: Levels of Alkaline Phosphatase

TBIL: Total Bilirubin

RBC: Red Blood Cells

HCT: Hematocrit

HGB: Hemoglobin

MCV: Mean Cell Volume

MCH: Mean Cell Hemoglobin

MCHC: Mean Corpuscular Hemoglobin Concentration

RETIC: Reticulocytes

WBC: White Blood Cells

NEU: Neutrophil

LYM: Lymphocytes

MONO: Monocytes

EOS: Eosinophil Count

BASO: Basophils

PLT: Number of Platelets

MPV: Mean Platelet Volume

## Introduction

Goats are important for both commercial and subsistent farming systems in rural southeastern United States. Limited resource producers keep goats primarily for meat and as a source of income since most subsistent farmers cannot afford to keep cattle. Goats are very versatile because not only do they provide goods but they also provide services such as vegetation management by eating unwanted vegetation in fields, and they can also prevent fuel fires by reducing fuel load [1]. The tremendous changes in demand that are ongoing within rural farm communities in the course of the last two decades has created the opportunity for goat production to become widespread [2]. Also, the substantial increase in the U.S. chevon imports in recent years, provided reasons for growth in the domestic meat-goat industry [3].

The Kiko goat is a composite breed developed in humid New Zealand in the 1970s and 1980s for improved growth and survivability by crossing Saanen, Toggenburg, and Nubian bucks with selected feral does. The Kiko is thought to be a vigorous, hardy, large frame, and early maturing animal that doesn't need pampering [4]. In the southeastern U.S., efficient meat goat production is difficult because warm, humid pasture conditions are optimum for gastrointestinal parasites. Internal parasites represent the greatest threat to goat productivity, health, and survival [5]. The Southeastern region is more favorable for Kiko breed of goats due to their increased parasitic resistance and lower incidence of hoof problems [6]. Moreover, the Kiko goat breed focuses heavily on the production of goat meat while maintaining minimal intervention inputs. Environmental conditions and in particular seasonal variations in air temperature and relative humidity are considered physiological stressors which affect the animal's biological system [7]. The variation in environmental variables such as ambient temperature, relative humidity, wind and rainfall were recognized as the potential hazards in livestock growth and production. In order to maintain homeothermic, an animal must be in thermal equilibrium with its environment, which includes radiation, air temperature, air movement and humidity [8].

Both serum biochemical and hematological profiles can be influenced by a number of factors such as age [9], physiological status [10, 11]. Biochemical and hematological parameters change in different seasons and temperature, humidity index may also be a factor in these fluctuations [12]. Reported elevations in lymphocytes, and phagocytic activities in Dhofari goat during the summer season, while a decrease in levels of neutrophils occurred in the winter season. [13] Reported that India Bengal goats express higher levels of hemoglobin, packed cell volume, mean corpuscular volume as well as mean corpuscular hemoglobin [14]. Reported that during the summer monsoon season, India Bengal goats showed higher levels of hemoglobin, packed cell volume, mean corpuscular volume as well as mean corpuscular hemoglobin [15,13], observed that ambient temperature and temperature humidity index have effects on both hematological and biochemical parameters in goats with total protein, blood urea nitrogen, and albumin significantly ( $P<0.01$ ) higher in summer than in both winter and autumn months [16]. Observed that in goats, summer season resulted in increases in levels of blood urea nitrogen, uric acid as well as creatinine. However other parameters such as calcium, cholesterol, glucose and inorganic phosphorus decreased during the summer months. While seasonal variations are extremely important to note, other environmental factors are equally important factors

when analyzing and determining hematological and biochemical parameters of meat goats breeds. It may be possible that blood values are also affected by both water temperature, and/or the differences between intensive and extensive production systems [17]. To date, very few hematological and serum biochemistry parameters for Kiko (meat) and their relationships to seasons of the year under semi-intensive management system have been published. The hematological and serum biochemical values obtained in this study will further underlined the need to establish appropriate physiological baseline values for Kiko meat goats in Alabama which could help in the realistic evaluation of management practice, nutrition, diagnosis of health as well as in determining the physiological status of goats used by limited resource producers. In addition, this study will provide insights into physiological responses of the Kiko meat goat to different seasons, allowing producers to better evaluate its ability to adapt and cope with various environmental stressors.

The objective of the present project was to investigate seasonal (spring, summer, fall, and winter) influences on the biochemical and hematological parameters of Kiko-meat goats managed under semi-intensive production system in southeast Alabama.

## Materials and Methods

### Animal management

This study was conducted at the Caprine Research and Education Unit of the George Washington Carver Agricultural Experiment Station at Tuskegee University, Tuskegee, Alabama (32.43N, 85.71W). Tuskegee is located in the southeastern region of the United States, sits 183 meters above sea level and has an annual precipitation amount of 1222mm. The Tuskegee University Animal Care and Use Committee approved the herd management protocol used.

in this project. For this project a total number of 15 Kiko goats that were semi-intensively managed in summer, fall, winter and spring were utilized. Goats were between the ages of 1 and 2 years old and were dewormed three times during the research period. All animals were managed on tall fescue (*Festuca arundinacea*) and Bermuda grass (*Cynodondactylon*) pastures and supplemented with Bermuda grass hay (*Cynodondactylon*) for ad libitum consumption. Animals were also supplemented with 341 g/d of alfalfa (17% crude protein, 1.5% crude fat, 30% crude fiber) and corn (7% crude protein, 3% crude fat, 4% crude fiber) and had access to trace mineral salt blocks. All research animals had access to water daily.

### Blood collection

Blood samples were collected between 8:00AM and 10:00AM once a week for three weeks for each of the calendar seasons, summer, fall, Winter and Spring. Blood samples were collected with an 18-gauge needle by jugular venipuncture. The blood was collected into two tubes, one in a plain tube for serum analysis and another with EDTA for whole blood analysis. A total of 5 mL of blood was collected per collection. Blood samples were centrifuged within two hours of collection and serum was harvested and stored at  $-20^{\circ}\text{C}$  for biochemical analysis. During each collection period temperature, humidity and rainfall were also recorded.

### Hematology

Samples for blood hematology was prepared by using a special blood analysis buffer approved for goat hematology (Concentrated

Lysing Reagent, SEACa and Florence, Italy). All samples were analyzed within 45 minutes after collection by using IDEXX Procyte Dx Automated Hematology Analyzer for total White Blood Cells (WBC), Red Blood Cells (RBCS) Hemoglobin HGB), hematocrit (HCT), Mean Corpuscular Volume (MCV), Mean Corpuscular Hemoglobin (MCH), Mean Corpuscular Hemoglobin Concentration (MCHC) and Number of Platelets (PLT). All samples were analyzed at the Tuskegee University College of veterinary medicine clinical pathology diagnostic laboratory.

### Serum biochemistry

Blood samples were left to clot and then centrifuged at 3000 rpm for 15 minutes, and then the serum was collected. The serum was kept frozen at -20°C until it was used for the biochemical analysis. The serum biochemistry was carried out using IDEXX Catalyst One Automated Chemistry Analyzer for Blood Urea Nitrogen (BUN) Aspartate aminotransferase (AST), Alanine Aminotransferase (ALT), Alkaline Phosphatase (ALP), Glucose (GLU), Creatinine (CREA), Calcium (CA) Total Protein (TP), and Total Bilirubin (TBIL). All samples were analyzed at the Tuskegee University College of veterinary medicine clinical pathology diagnostic laboratory.

### Body weight and body condition score

Body weight was recorded using a digital scale. In addition to assessing body condition, a palpable body condition scoring scale was also used on a scale of 1 to 5, (1 being emaciated and 5 being obese). Body condition scoring was subjectively determined. Body weight was taken each time blood collections occurred for consistent values.

### Environmental conditions

The mean values of daily and cumulative rainfall, rainy days, relative humidity, and minimum, maximum temperature and average summer fall, winter and spring seasons are shown in (Table 1.)

| Item                         | Season | Mean  | SD   | Minimum | Maximum |
|------------------------------|--------|-------|------|---------|---------|
| Temperature at Collection °C | Spring | 13.3  | 2.55 | 11.1    | 16.1    |
|                              | Summer | 23.6  | 2.50 | 21.1    | 26.1    |
|                              | Fall   | 13.8  | 8.61 | 8.88    | 23.8    |
|                              | Winter | 6.08  | 11.2 | -2.7    | 18.8    |
| High °C                      | Spring | 25.7  | 3.59 | 21.6    | 28.3    |
|                              | Summer | 32.9  | 1.15 | 31.6    | 33.8    |
|                              | Fall   | 23.6  | 1.27 | 22.2    | 24.4    |
|                              | Winter | 16.07 | 7.85 | 7.22    | 22.2    |
| Low °C                       | Spring | 10.1  | 0.28 | 10.0    | 10.5    |
|                              | Summer | 21.1  | 1.61 | 19.4    | 22.2    |
|                              | Fall   | 6.47  | 2.31 | 3.88    | 8.33    |
|                              | Winter | 4.98  | 10.5 | -3.88   | 16.6    |
| Humidity%                    | Spring | 71.6  | 18.1 | 55.0    | 91.0    |
|                              | Summer | 83.3  | 7.63 | 75.0    | 90.0    |
|                              | Fall   | 88.0  | 8.0  | 80.0    | 96.0    |
|                              | Winter | 82.0  | 21.6 | 57.0    | 95.0    |
| Rainfall cm                  | Spring | 0.00  | 0.00 | 0.00    | 0.00    |
|                              | Summer | 0.02  | 0.04 | 0.00    | 0.07    |
|                              | Fall   | 0.00  | 0.00 | 0.00    | 0.00    |
|                              | Winter | 0.87  | 1.51 | 0.00    | 2.61    |

**Table 1:** Environmental Conditions.

### Statistical analysis

All results are expressed as ± Standard Deviation (SD) [18]. Statistical software calculated the minimum and maximum values to determine the range, mean, and standard deviation of the mean. The effects of season on biochemical parameters and hematological profiles of goats were Analyzed by an Analysis of Variance (ANOVA) using [18], statistical software.

### Results and discussion

#### Body weight, body condition score and serum biochemical parameters

Body weight of 59.9 kg ± 11.7 kg was recorded for spring, 46.2 kg ± 11.4 kg in summer, 46.3 kg ± 13.6 kg, in fall, and 46.5 kg ± 10.1 kg in winter respectively. For body condition scores, results show 3.0 ± 0.0 in the spring, 2.7 ± 0.4 in the summer, 2.6 ± 0.5 in the fall, and 2.6 ± 0.5 in the winter respectively. Seasonal values for the following serum biochemical parameters GLU = Glucose, CREA = Creatinine, BUN = Blood Urea Nitrogen, CA = Calcium TP = Total Protein, ALT= Alkaline Aminotransferase, AST= Aspartate Aminotransferase, ALKP = Levels of Alkaline Phosphatase, TBIL = Total Bilirubin are presented in (Table 2).

| Item                         | Season | Mean  | SD   | Minimum | Maximum |
|------------------------------|--------|-------|------|---------|---------|
| Temperature at Collection °C | Spring | 13.3  | 2.55 | 11.1    | 16.1    |
|                              | Summer | 23.6  | 2.50 | 21.1    | 26.1    |
|                              | Fall   | 13.8  | 8.61 | 8.88    | 23.8    |
|                              | Winter | 6.08  | 11.2 | -2.7    | 18.8    |
| High °C                      | Spring | 25.7  | 3.59 | 21.6    | 28.3    |
|                              | Summer | 32.9  | 1.15 | 31.6    | 33.8    |
|                              | Fall   | 23.6  | 1.27 | 22.2    | 24.4    |
|                              | Winter | 16.07 | 7.85 | 7.22    | 22.2    |
| Low °C                       | Spring | 10.1  | 0.28 | 10.0    | 10.5    |
|                              | Summer | 21.1  | 1.61 | 19.4    | 22.2    |
|                              | Fall   | 6.47  | 2.31 | 3.88    | 8.33    |
|                              | Winter | 4.98  | 10.5 | -3.88   | 16.6    |
| Humidity%                    | Spring | 71.6  | 18.1 | 55.0    | 91.0    |
|                              | Summer | 83.3  | 7.63 | 75.0    | 90.0    |
|                              | Fall   | 88.0  | 8.0  | 80.0    | 96.0    |
|                              | Winter | 82.0  | 21.6 | 57.0    | 95.0    |
| Rainfall cm                  | Spring | 0.00  | 0.00 | 0.00    | 0.00    |
|                              | Summer | 0.02  | 0.04 | 0.00    | 0.07    |
|                              | Fall   | 0.00  | 0.00 | 0.00    | 0.00    |
|                              | Winter | 0.87  | 1.51 | 0.00    | 2.61    |

**Table 2:** Descriptive statistics for seasonal serum biochemical profiles 1 of kiko meatgoats.

1GLU = Glucose, CREA = Creatinine, BUN = Blood Urea Nitrogen, CA = Calcium TP = Total Protein, ALT= Alkaline Aminotransferase, AST= Aspartate Aminotransferase, ALKP = Levels of Alkaline Phosphatase, TBIL = Total Bilirubin

These results were compared with values obtained from other goat breeds [19] and the reference range for goats as reported by [20]. GLU levels of 65.0 ± 14.9, 62.7 ± 6.2, 60.6 ± 16.6 and 57.8 ± 11.3 mg/DL were obtained for spring, summer, fall and winter seasons

respectively. CREA values were  $0.52 \pm 0.07$ ,  $0.54 \pm 0.15$ ,  $0.62 \pm 0.16$ ,  $0.63 \pm 0.09$  mg/DL for spring, summer, fall and winter seasons respectively. Mean values of  $15.6 \pm 4.18$ ,  $15.8 \pm 5.17$ ,  $12.4 \pm 3.7$  and  $9.3 \pm 4.02$  mg/DL for BUN in spring, summer, fall and winter seasons respectively. Lower values of TBIL, BUN were found in Kiko goat breeds when compared to values from other goat breeds [21]. Lower concentration of BUN is an indication of the dietary protein level or liver chronic diseases [22]. Mean values of  $8.2 \pm 0.6$ ,  $9.1 \pm 0.6$ ,  $8.4 \pm 1.1$  and  $8.4 \pm 0.5$  mg/DL were recorded for CA in spring, summer, fall and winter seasons respectively. TP values of  $7.1 \pm 0.5$ ,  $7.8 \pm 0.7$ ,  $7.6 \pm 0.9$  and  $7.8 \pm 0.5$  g/Dl were obtained in spring, summer, fall and winter seasons respectively. In addition, levels of ALKP were  $90.8 \pm 58.3$ ,  $97.9 \pm 59.3$ ,  $47.8 \pm 17.1$  and  $80.0 \pm 30.7$  in spring, summer, fall and winter seasons respectively. TBIL, AST, GLU and CREA levels were not significantly affected by seasons ( $P \geq 0.05$ ). CA and BUN levels were highly significantly influenced by seasons ( $P \leq 0.01$ ). Total protein and alkaline phosphatase levels showed significant seasonal variations ( $P \leq 0.05$ ) as shown in (Table 3).

| TRAIT      | SEASON          |                 |                 |                 | P-values | Significance |
|------------|-----------------|-----------------|-----------------|-----------------|----------|--------------|
|            | SPRING          | SUMMER          | FALL            | WINTER          |          |              |
| GLU mg/DL  | $65.0 \pm 14.9$ | $62.7 \pm 6.2$  | $60.6 \pm 16.6$ | $57.8 \pm 11.8$ | 0.49     | NS           |
| CREA mg/DL | $0.52 \pm 0.07$ | $0.54 \pm 0.15$ | $0.62 \pm 0.16$ | $0.63 \pm 0.09$ | 0.06     | NS           |
| BUN mg/DL  | $15.6 \pm 4.1$  | $15.8 \pm 5.1$  | $12.4 \pm 3.7$  | $9.3 \pm 4.02$  | 0.009    | **           |
| BUN/CREA   | $30.6 \pm 10.6$ | $30.8 \pm 11.0$ | $21.6 \pm 8.1$  | $15.8 \pm 6.6$  | 0.001    | **           |
| CA mg/DL   | $8.2 \pm 0.6$   | $9.1 \pm 0.6$   | $8.4 \pm 1.1$   | $8.4 \pm 0.5$   | 0.008    | **           |
| TP g/Dl    | $7.1 \pm 0.5$   | $7.8 \pm 0.7$   | $7.6 \pm 0.9$   | $7.8 \pm 0.5$   | 0.016    | *            |
| ALT U/L    | $11.6 \pm 2.1$  | $19.2 \pm 6.2$  | $12.8 \pm 5.6$  | $11.2 \pm 3.6$  | 0.001    | **           |
| AST U/L    | $99.6 \pm 30.7$ | $87.5 \pm 18.4$ | $78.6 \pm 11.8$ | $83.2 \pm 28.0$ | 0.94     | NS           |
| ALKP U/L   | $90.8 \pm 58.3$ | $97.9 \pm 59.3$ | $47.8 \pm 17.1$ | $80.0 \pm 30.7$ | 0.018    | *            |
| TBIL mg/Dl | $0.46 \pm 0.15$ | $0.50 \pm 0.09$ | $0.38 \pm 0.22$ | $0.39 \pm 0.21$ | 0.210    | NS           |

**Table 3:** Seasonal differences in serum biochemical profiles of kiko meat goats (p- values).

\*Significant if  $P \leq 0.05$ ,\*\* Highly Significant if  $P \leq 0.01$ ,\*\*\*NS = Not Significant

Our study showed a moderate increase in total protein and globulin which is associated with the rise in ambient summer temperature [23]. Suggested that any such increase in serum protein could be a physiological attempt to maintain extended plasma volume. Variations in serum protein concentration were observed in lactating cattle and buffaloes during spring and summer seasons [24]. The values of creatinine in the current study did not increase under high environmental temperatures in summer suggesting that the Kiko meat goats are metabolically adapted to hot and humid conditions of southeast Alabama. In the present study blood urea values were higher at highest temperatures in summer months, confirming results obtained [25], could be attributed to an increased utilization of amino acids as energy source in hot humid conditions.

Determination of serum total bilirubin is valuable for the diagnosis of fatty liver in ruminants. Total bilirubin is often increased after parturition [26] and is also increased during periods of anorexia [27]. The results of our study showed lower ALP values than that reported in Sokoto red, Sahel and Saanen goats [28]. However, our ALP values were slightly higher than that reported in West African Dwarf (WAD) goats [29]. ALP constitutes a large group of isoenzymes, which plays important roles in the transportation of sugar, phosphate and it originates from different tissues such as liver, bone, placenta, and intestine [28].

### Hematology

Mean values and standard deviations for RBC = Red Blood Cells, HCT= Hematocrit, HGB = Hemoglobin, MCV = Mean Cell Volume, MCH = Mean Cell Hemoglobin, MCHC = Mean Corpuscular Hemoglobin Concentration, RETIC= Reticulocytes, WBC = White Blood Cells, NEU = Neutrophil, LYM = Lymphocytes, MONO = Monocytes, EOS = Eosinophil Count, BASO= Basophils, PLT= Number of Platelets, MPV = Mean Platelet Volume in the summer, fall, winter and spring seasons are shown in (table 4). The average values obtained in this study for RBC, PCV, HGB, MCV, MCH, MCHC are within the reference ranges for the caprine species (table 4). The mean RBC varied in all seasons from a mean low count of  $16.7 \pm 4.12$  M/ $\mu$ l in spring to a mean high count of  $20.5 \pm 4.27$  M/ $\mu$ l in winter. These differences were significant ( $P \leq 0.05$ ) (table 5). Also, values of RBC, HGB, and HCT in our study were within the normal range for goats as reported by [30], similar to those observed in Barbari Black Aardi breed of goats [31]. In contrast, the values of RBCs, HGB, and HCT were higher than that reported in the Damascus breed [30]. Higher values of MCV and MCH and lower values of MCHC were observed in the Kiko goat breed comparing to that in the other goat breeds [28], as shown in (Table 4).

| Item             | Season | Mean | Standard Deviation | Minimum | Maximum | Reference Value |
|------------------|--------|------|--------------------|---------|---------|-----------------|
| RBC( M/ $\mu$ l) | Spring | 16.7 | 4.12               | 12.5    | 23.5    |                 |
|                  | Summer | 18.1 | 2.52               | 14.3    | 22.2    |                 |
|                  | Fall   | 18.0 | 1.58               | 15.6    | 21.3    |                 |
|                  | Winter | 20.5 | 4.27               | 16.1    | 28.0    |                 |
|                  |        |      |                    |         |         | 10.32-23.43     |

|                  |        |      |      |       |      |           |
|------------------|--------|------|------|-------|------|-----------|
| HCT %            | Spring | 36.1 | 7.3  | 23.0  | 45.4 |           |
|                  | Summer | 29.0 | 9.8  | 16.0  | 46.7 |           |
|                  | Fall   | 27.2 | 9.4  | 13.7  | 41.4 |           |
|                  | Winter | 32.3 | 11.3 | 11.1  | 51.4 |           |
|                  |        |      |      |       |      | 22.0-39.0 |
| HGB G/ DL        | Spring | 9.6  | 0.75 | 8.5   | 10.9 |           |
|                  | Summer | 9.2  | 1.4  | 7.1   | 11.5 |           |
|                  | Fall   | 8.8  | 1.2  | 6.5   | 10.8 |           |
|                  | Winter | 9.9  | 1.5  | 8.2   | 13.7 |           |
|                  |        |      |      |       |      | 8.9-13.8  |
| MCV fL           | Spring | 23.5 | 8.7  | 10.9  | 32.6 |           |
|                  | Summer | 16.2 | 5.3  | 7.6   | 24.4 |           |
|                  | Fall   | 15.0 | 4.9  | 8.3   | 20.9 |           |
|                  | Winter | 16.3 | 5.7  | 4.9   | 23.7 |           |
|                  |        |      |      |       |      | 14.0-22.3 |
| MCH pg           | Spring | 6.0  | 1.0  | 4.4   | 7.3  |           |
|                  | Summer | 5.1  | 0.64 | 4.1   | 6.2  |           |
|                  | Fall   | 4.8  | 0.58 | 4.0   | 5.7  |           |
|                  | Winter | 4.9  | 0.76 | 3.8   | 6.2  |           |
|                  |        |      |      |       |      | 5.0-7.0   |
| MCHC g/dL        | Spring | 28.2 | 8.1  | 21.6  | 40.9 |           |
|                  | Summer | 34.3 | 9.8  | 24.6  | 53.8 |           |
|                  | Fall   | 35.2 | 9.0  | 26.1  | 48.9 |           |
|                  | Winter | 35.1 | 15.4 | 20.2  | 78.4 |           |
|                  |        |      |      |       |      | 32.0-34.0 |
| RETIC K/ $\mu$ L | Spring | 0.01 | 0.03 | 0.00  | 0.1  |           |
|                  | Summer | 0.00 | 0.00 | 0.00  | 0.00 |           |
|                  | Fall   | 0.00 | 0.00 | 0.00  | 0.00 |           |
|                  | Winter | 0.00 | 0.00 | 0.00  | 0.00 |           |
|                  |        |      |      |       |      | 0.0-15.0  |
| WBC K/ $\mu$ L   | Spring | 15.0 | 3.5  | 10.7  | 25.7 |           |
|                  | Summer | 15.9 | 6.4  | 7.9   | 28.3 |           |
|                  | Fall   | 14.3 | 5.0  | 7.4   | 22.0 |           |
|                  | Winter | 14.8 | 6.5  | 7.1   | 29.3 |           |
|                  |        |      |      |       |      | N/A       |
| NEU %            | Spring | 53.8 | 15.2 | 26.9  | 71.2 |           |
|                  | Summer | 39.7 | 11.1 | 22.2  | 58.7 |           |
|                  | Fall   | 44.5 | 5.1  | 32.71 | 58.2 |           |
|                  | Winter | 42.5 | 7.9  | 30.9  | 55.4 |           |
|                  |        |      |      |       |      | N/A       |
| LYM %            | Spring | 36.7 | 10.9 | 25.7  | 58.8 |           |
|                  | Summer | 44.3 | 6.8  | 31.6  | 56.6 |           |
|                  | Fall   | 37.7 | 5.8  | 23.8  | 47.9 |           |
|                  | Winter | 41.4 | 5.1  | 34.9  | 54.3 |           |
|                  |        |      |      |       |      | N/A       |
| MONO %           | Spring | 4.4  | 3.5  | 0.2   | 9.9  |           |
|                  | Summer | 7.8  | 4.5  | 1.7   | 16.1 |           |
|                  | Fall   | 8.9  | 3.2  | 3.8   | 15.7 |           |
|                  | Winter | 7.7  | 3.5  | 3.3   | 14.9 |           |
|                  |        |      |      |       |      | N/A       |

|                 |        |       |       |       |        |            |
|-----------------|--------|-------|-------|-------|--------|------------|
| EOS %           | Spring | 4.1   | 2.8   | 0.6   | 8.7    |            |
|                 | Summer | 7.6   | 4.3   | 2.7   | 14.8   |            |
|                 | Fall   | 8.5   | 5.4   | 1.2   | 19.4   |            |
|                 | Winter | 7.7   | 3.9   | 1.4   | 14.7   |            |
|                 |        |       |       |       |        | N/A        |
| BASO %          | Spring | 0.75  | 0.35  | 0.1   | 1.2    |            |
|                 | Summer | 0.42  | 0.28  | 0.1   | 1.10   |            |
|                 | Fall   | 0.24  | 0.16  | 0.00  | 0.60   |            |
|                 | Winter | 0.42  | 0.28  | 0.10  | 1.10   |            |
|                 |        |       |       |       |        | N/A        |
| NEU K/ $\mu$ L  | Spring | 8.0   | 2.9   | 4.2   | 16.5   |            |
|                 | Summer | 5.9   | 1.9   | 3.3   | 8.9    |            |
|                 | Fall   | 6.3   | 2.6   | 2.7   | 10.9   |            |
|                 | Winter | 6.2   | 3.3   | 3.1   | 16.2   |            |
|                 |        |       |       |       |        | 1.72-10.61 |
| LYM K/ $\mu$ L  | Spring | 5.5   | 2.02  | 3.2   | 9.3    |            |
|                 | Summer | 7.2   | 3.1   | 2.5   | 12.4   |            |
|                 | Fall   | 5.3   | 1.8   | 2.9   | 7.7    |            |
|                 | Winter | 6.2   | 3.1   | 2.7   | 12.7   |            |
|                 |        |       |       |       |        | 2.68-11.54 |
| MONO K/ $\mu$ L | Spring | 0.63  | 0.50  | 0.04  | 1.4    |            |
|                 | Summer | 1.3   | 1.04  | 0.20  | 3.2    |            |
|                 | Fall   | 1.30  | 0.65  | 0.36  | 2.3    |            |
|                 | Winter | 1.1   | 0.68  | 0.34  | 2.1    |            |
|                 |        |       |       |       |        | 0.06-0.89  |
| EOS K/ $\mu$ L  | Spring | 0.68  | 0.58  | 0.08  | 2.05   |            |
|                 | Summer | 1.4   | 1.3   | 0.29  | 4.1    |            |
|                 | Fall   | 1.2   | 1.08  | 0.22  | 4.10   |            |
|                 | Winter | 1.08  | 0.76  | 0.24  | 3.5    |            |
|                 |        |       |       |       |        | 0.03-1.29  |
| BASO K/ $\mu$ L | Spring | 0.11  | 0.07  | 0.01  | 0.32   |            |
|                 | Summer | 0.06  | 0.03  | 0.02  | 0.15   |            |
|                 | Fall   | 0.03  | 0.02  | 0.00  | 0.07   |            |
|                 | Winter | 0.06  | 0.04  | 0.02  | 0.17   |            |
|                 |        |       |       |       |        | 0.00-0.24  |
| PLT K/ $\mu$ L  | Spring | 641.6 | 167.4 | 322.0 | 906.0  |            |
|                 | Summer | 591.4 | 143.7 | 329.0 | 822.0  |            |
|                 | Fall   | 749.6 | 248.4 | 527.0 | 1517.0 |            |
|                 | Winter | 645.5 | 249.5 | 136.0 | 1116.0 |            |
|                 |        |       |       |       |        | 246-912    |
| MPV fL          | Spring | 7.8   | 0.3   | 7.1   | 8.3    |            |
|                 | Summer | 7.8   | 0.3   | 7.1   | 8.3    |            |
|                 | Fall   | 7.9   | 0.4   | 7.0   | 8.4    |            |
|                 | Winter | 8.2   | 0.3   | 7.6   | 8.7    |            |
|                 |        |       |       |       |        | N/A        |

**Table 4:** Descriptive statistics for seasonal hematological profiles of kiko meat goats.

1 RBC = Red Blood Cells, HCT= Hematocrit, HGB = Hemoglobin, MCV = Mean Cell Volume,

MCH = Mean Cell Hemoglobin, MCHC = Mean Corpuscular Hemoglobin Concentration, RETIC= Reticulocytes, WBC = White Blood Cells, NEU = Neutrophil, LYM = Lymphocytes, MONO = Monocytes, EOS = Eosinophil Count, BASO= Basophils, PLT= Number of Platelets, MPV = Mean Platelet Volume.



| TRAIT            | SEASON            |                   |                   |                   | P-values | Significance |
|------------------|-------------------|-------------------|-------------------|-------------------|----------|--------------|
|                  | SPRING            | SUMMER            | FALL              | WINTER            |          |              |
| RBC (M/ $\mu$ L) | 16.7 $\pm$ 4.12   | 18.1 $\pm$ 2.52   | 18.0 $\pm$ 1.58   | 20.5 $\pm$ 4.27   | 0.0269   | *            |
| HCT %            | 36.1 $\pm$ 7.39   | 29.0 $\pm$ 9.86   | 27.2 $\pm$ 9.43   | 32.3 $\pm$ 11.3   | 0.0671   | NS           |
| HGB G/DL         | 9.67 $\pm$ 0.75   | 9.23 $\pm$ 1.47   | 8.82 $\pm$ 1.21   | 9.99 $\pm$ 1.59   | 0.0870   | NS           |
| MCV fL           | 23.5 $\pm$ 8.72   | 16.2 $\pm$ 5.37   | 15.0 $\pm$ 4.97   | 16.3 $\pm$ 5.76   | 0.0022   | **           |
| MCH pg           | 6.00 $\pm$ 1.03   | 5.10 $\pm$ 0.64   | 4.87 $\pm$ 0.58   | 4.96 $\pm$ 0.76   | 0.0006   | **           |
| MCHC g/dL        | 28.6 $\pm$ 8.12   | 34.3 $\pm$ 9.82   | 35.2 $\pm$ 9.07   | 35.1 $\pm$ 15.4   | 0.2559   | NS           |
| RETIC K/ $\mu$ L | 3.76 $\pm$ 2.29   | 1.34 $\pm$ 2.02   | 0.90 $\pm$ 1.26   | 1.56 $\pm$ 1.49   | 0.0003   | **           |
| WBC K/ $\mu$ L   | 15.04 $\pm$ 3.53  | 15.9 $\pm$ 6.47   | 14.3 $\pm$ 5.02   | 14.8 $\pm$ 6.59   | 0.8694   | NS           |
| NEU %            | 8.05 $\pm$ 2.99   | 5.94 $\pm$ 1.98   | 6.38 $\pm$ 2.60   | 6.27 $\pm$ 3.31   | 0.1678   | NS           |
| LYM %            | 36.7 $\pm$ 10.9   | 44.3 $\pm$ 6.84   | 37.7 $\pm$ 5.87   | 41.4 $\pm$ 5.17   | 0.0294   | *            |
| MONO %           | 4.48 $\pm$ 3.50   | 7.82 $\pm$ 4.54   | 8.91 $\pm$ 3.29   | 7.74 $\pm$ 3.54   | 0.0135   | *            |
| EOS %            | 0.68 $\pm$ 0.58   | 1.41 $\pm$ 1.33   | 1.25 $\pm$ 1.08   | 1.08 $\pm$ 0.76   | 0.2229   | NS           |
| BASO %           | 0.70 $\pm$ 0.35   | 0.42 $\pm$ 0.28   | 0.24 $\pm$ 0.16   | 0.44 $\pm$ 0.23   | 0.0002   | **           |
| LYM K/ $\mu$ L   | 5.54 $\pm$ 2.02   | 7.22 $\pm$ 3.15   | 5.32 $\pm$ 1.86   | 6.26 $\pm$ 3.12   | 0.1973   | NS           |
| MONO K/ $\mu$ L  | 0.63 $\pm$ 0.50   | 1.34 $\pm$ 1.04   | 1.30 $\pm$ 0.65   | 1.15 $\pm$ 0.68   | 0.448    | NS           |
| EOS K/ $\mu$ L   | 0.68 $\pm$ 0.58   | 1.41 $\pm$ 1.33   | 1.25 $\pm$ 1.08   | 1.08 $\pm$ 0.76   | 0.2229   | NS           |
| BASO K/ $\mu$ L  | 0.11 $\pm$ 0.07   | 0.06 $\pm$ 0.03   | 0.03 $\pm$ 0.02   | 0.06 $\pm$ 0.04   | 0.0009   | **           |
| PLT K/ $\mu$ L   | 641.6 $\pm$ 167.4 | 591.4 $\pm$ 143.7 | 749.6 $\pm$ 248.1 | 645.5 $\pm$ 248.5 | 0.2142   | NS           |
| MPV fL           | 7.81 $\pm$ 0.35   | 7.80 $\pm$ 0.33   | 7.92 $\pm$ 0.43   | 8.22 $\pm$ 0.35   | 0.0090   | **           |

**Table 5:** Seasonal differences in hematological profiles 1 of kiko meat goats (p-values)

\*Significant if  $P \leq 0.05$ , \*\* Highly Significant if  $P \leq 0.01$ , \*\*\*NS = Not Significant

1 RBC = Red Blood Cells, HCT=Hematocrit, HGB = Hemoglobin, MCV =Mean Cell Volume, MCH = Mean Cell Hemoglobin, MCHC = Mean Corpuscular Hemoglobin Concentration, RETIC= Reticulocytes, WBC =22 White Blood Cells, NEU = Neutrophils, LYM = Lymphocytes, MONO = Monocytes, EOS = Eosinophil Count,BASO= Basophils, PLT= Number of Platelets MPV = Mean Platelet Volume

As shown in (tables 4&5) the mean HGB concentration decreased slightly between winter and spring ( $9.9 \pm 1.5$  vs.  $9.6 \pm 0.75$  G/DL), however, no significant seasonal differences were observed ( $P \leq 0.05$ ). Between spring and winter, the mean HCT increased slightly ( $36.1 \pm 7.39$  vs.  $32.3 \pm 11.3\%$ ). Intermediate values were found in fall and summer ( $27.2 \pm 9.43$  vs.  $29.0 \pm 9.86\%$ ) respectively. These differences were not significant ( $P \geq 0.05$ ). The Mean Corpuscular Hemoglobin Concentration (MCHC) has a reference value of 32.0-34.0 g/dL for goats [30]. MCHC increased from  $28.6 \pm 8.12$  g/dL in spring to reach the highest value of  $35.2 \pm 9.07$  g/dL in the fall. There were fluctuating variations throughout all four seasons thus indicating seasonal influence, however, the values obtained may represent possible iron deficiency in diets consumed by these goats. The decrease levels of the MCHC, especially during the fall and winter season, may also be indicative of a parasitic infection.

Highly significant seasonal differences ( $P \leq 0.01$ ) were observed for both mean corpuscular volume and mean cell hemoglobin ( $23.5 \pm 8.72$ ,  $16.2 \pm 5.37$ ,  $15.0 \pm 4.97$  and  $16.3 \pm 5.76$  fl) ( $6.00 \pm 1.03$ ,  $5.10 \pm 0.64$ ,  $4.87 \pm 0.58$  and  $4.96 \pm 0.76$  pg for spring, summer fall and winter respectively. Results from the current show that the (MCV) and (MCH) were significantly lower in winter compared to values obtained in spring. The lowest MCV ( $15.0 \pm 4.97$ fl) was obtained in fall. This low (MCV) value could be related to the negative correlation between size and number of erythrocytes as suggested [32]. The values obtained in the present study for (MCV) and (MCH) during winter and wet summer were not in general agreement with other findings [33], who reported high values of (MCV) and (MCH) in cold dry environment compared to values in hot humid conditions. Non-significant seasonal changes ( $P \geq 0.05$ ) in total leucocyte count (WBC) were found ( $15.04 \pm 3.53$ ,  $15.9 \pm 6.47$ ,  $14.3 \pm 5.02$  and  $14.8 \pm 6.59$  K/ $\mu$ L) for spring, summer, fall and winter respectively. The present results for seasonal changes in erythrocyte count and (HGB) concentration in Kiko goats are in agreement with the findings in other breeds of goats [33,34], reported higher values of erythrocyte count, (PCV) and (HGB) during summer compared to winter months. These variations between Kiko and other breeds of goats may be attributed to differences in environmental conditions as well as nutritional factors [35]. Reported lower values of these indices in winter for Cameroon goats kept in temperate environment and attributed that to the change in diet. The effect that seasonality can have on WBC count is attributed, among other factors to the stress associated with the cold in winter, leading to suppression of the immune response [34,36], reported that elevation of WBC in winter occurs because the lymphoid organs tend to become larger in colder seasons. Our results show that the Kiko goats had higher WBCs and neutrophils percentage compared to Babari, Black Aardi, Damascus, Kano brown and Nigerian Sahel goats [15]. Also, the WBCs in the Kiko goats were higher than the reference range as reported by [30].

RETIC values for all four seasons remained within the reference value range of 0.0-15.0 K/ $\mu$ L. The only fluctuation occurred during the spring with  $0.01 \pm 0.03$  and a minimum value of 0.00 K/ $\mu$ L and with a maximum value of 0.1 K/ $\mu$ L. Neutrophils percentage (NEU) was higher during the spring ( $53.8 \pm 15.2$ ) compared to summer ( $39.7 \pm 11.1$ ), fall ( $44.5 \pm 5.1$ ), and winter ( $42.5 \pm 7.9$ ). Basophil K/ $\mu$ L (BASO) had no seasonal fluctuations aside from exceeding the reference value (0.00-2.24 K/ $\mu$ L) during spring (0.32 K/ $\mu$ L). Number of platelets (PLT) K/ $\mu$ L, has a reference value of 246-914 K/ $\mu$ L, no seasonal increases above this reference range were recorded. While the mean platelet volume (MPV) has no particular reference value

for goats, values obtained in this study were significantly influenced by seasons ( $7.81 \pm 0.35$ ,  $7.80 \pm 0.33$ ,  $7.92 \pm 0.43$ , and  $8.22 \pm 0.35$  fL;  $P \leq 0.05$ ). The percentage of monocytes, eosinophils and basophils in this study agreed with the data reported on the Barbari and black Aardi breed of goats [10] but slightly lower than the reported value in Damascus breed of goats [37]. However, these values were within the normal range reported by [30].

In this present study hematological and serum biochemical parameters were compared to standard, laboratory-dependent reference values. These reference values generally represent a 95% confidence interval. This means that 95% of normal animals should have a given blood profiles within this range. This also suggests that 5% of the population will be outside of this reference range and still be normal, emphasizing the need to clinically evaluate the animal. A number of factors, most notably physiologic state and age have been shown to influence blood hematological and serum biochemical parameters. Most reference ranges do not account for these differences and thus may confound direct interpretation. A new approach to hematological and serum biochemical profiling, which involves pooling larger sample numbers, specific animal selection relative to physiologic state and stage of production, must be examined in an effort to better interpret blood metabolite concentrations on a herd basis. Most importantly it must be remembered that metabolic profiles are almost useless without being coupled with environment and facility evaluations, body condition scoring and diet evaluation. It is only when the whole picture is evaluated will the uses of hematological and serum biochemical profiles produce useful diagnostic information.

## Conclusion and Recommendations

The results of the present study suggest and show that seasonal variations do have a role on hematological and serum biochemical profiles in Kiko meat goats. The higher red blood cell values observed in this study may likely be a sign of healthier goats. The components of some of the hematological and serum biochemical parameters in Kiko goats in this study seem to point out some differences from those obtained for other meat goat breeds. Furthermore, the observed differences further support the fact that the physiological parameters reported for other meat goat breeds may not be applied on Kiko goats kept in this ecological zone (southeast Alabama). It is concluded that hematological and serum biochemical parameters were mostly within the physiological range for goats as reported from other studies. This data can contribute to our knowledge for monitoring health status, diagnosis of disease and management in this breed in Alabama. Moreover, this study underscores the need to establish appropriate physiological baseline values for meat breed goats in Alabama which could help in realistic valuation of the management practice, nutrition, diagnosis of health as well as in determining the physiological status of goats among limited resource producers who often prefer the Kiko breed of meat goat.

## Acknowledgments

The authors wish to acknowledge the faculty and staff of Caprine Research Unit, George

Washington Carver Experimental Station, College of Agriculture, Environmental & Nutrition

Sciences (CAENS), Tuskegee University for technical support and USDA-NIFA for financial support.

## References

1. Solaiman SG (2007) Assessment of the meat goat industry and future outlook for US small farms. Tuskegee University.
2. Hart SP (2001) Recent perspectives in using goats for vegetation management in the USA. *Journal of Dairy Science*. 84: 170-176.
3. Sande DN, Houston JE, Epperson JE (2005) The relationship of consuming populations to meat-goat production in the United States. *J. Food Distribution Research* 36:156-160.
4. Browning L (2006) Breed Options for Meat Goat Production in Alabama. Alabama Cooperative Extension System Alabama A and M University and Auburn University Pub UNP-0084.
5. Kaplan RM, Burke JM, Terrill TH, Miller JE, Getz WR, et al. (2004) Validation of the FAMACHA® eye color chart for detecting clinical anemia on sheep and goat farms in the southern United States. *Vete Paras* 123: 105-120.
6. Wade K (2004) Kiko goats in the USA. *New Zealand Rare Breeds New*.
7. El-Nouty FD, Al-Haidary AA, Salah MS (1990) Seasonal variation in hematological values of high-and average yielding Holstein cattle in semi-arid environment. *J King Saud Univ* 2: 173-182.
8. Kadzere CT, Murphy MR, Silanikove N, Maltz E (2002) Heat stress in lactating dairy cows a review. *Livestock Prod Sci* 77: 59-91.
9. Oduye O (1976) Hematological values in Nigeria goats and sheep. *Tropical Animal Health and Production*, 8: 131-136.
10. Piccione G, Borruso M, Fazio F, Giannetto C, Caola G (2007) Physiological parameters in lambs during the first 30 days postpartum. *Small Rumin. Res* 72: 57-60.
11. Shehab-El-Deen MA, Fadel MS, van Soom A, Saleh SY, Maes D, et al. (2010) Circadian rhythm of metabolic changes associated with summer heat stress in high-producing dairy cattle. *Trop Anim Health Pro* 42: 1119-112.
12. Ribeiro MN, Ribeiro NL, Bozzi R, Costa RG (2018) Physiological and biochemical blood variables of goats subjected to heat stress a review *J of Applied Ani Res* 46: 1036-1041.
13. Bhatta M, Das D, Ranjan P (2014) The Effect of Ambient Temperature on Some Biochemical Profiles of Black Bengal Goats *Capra aegagrus hircus* In Two Different Agro-Climatic Zones in West Bengal India. *J of Pharm and Bio Sci* 9: 32-36.
14. Banerjee D, Upadhyay RC, Chaudhary UB, Kumar R, Singh S, et al. (2015) Seasonal variations in physio-biochemical profiles of Indian goats in the paradigm of hot and cold climate. *Biological rhythm research* 46: 221-236.
15. Arfuso F, Fazio F, Rizzo M, Marafioti S, Zanghi E, et al. (2016) Factors affecting the hematological parameters in different goat breeds from Italy. *Ann Anim Sci* 16: 743-757.
16. Silanikove N (2000). The physiological basis of adaptation in goats to harsh environments. *Small Ruminant Res* 35: 181-193.
17. Maloiy GMO, Taylor CR (1971) Water requirements of African goats and haired sheep. *J Agri Sci Cambridge* 77: 203-208.
18. (1996) SAS/STAT1 Software. Changes and Enhancements through Release .
19. Tambuwai F, Agaie B, Bangana A (2002). Haematological and Biochemical Values of Apparently Healthy Red Sokoto Goats. *Proceedings of the 27 Annual Conference of the Nigerian Soci for Ani Prod* 50-53.
20. Latimer, Duncan (2011) Prasse's Veterinary Laboratory Medicine: Clinical Pathology. Ames IA: Wiley-Blackwell.
21. Egbe-Nwiyi T, Igwenagu E, Samson M (2015) The influence of sex on the haematological values of apparently healthy adult Nigerian Sahel goats. *Sokoto J Vet Sci* 13: 54-58.

22. Mishra A, Chatterjee US, Mandal TK (2013) Induction of chronic renal failure in goats using Cisplatin: A new animal model. *Toxicol Int* 20: 56-60.
23. Ganaie AH, Ghasura RS, Mir NA, Bumla NA, Sankar G, et al. (2013) Biochemical and Physiological Changes during Thermal Stress in Bovines: A Review. *Iranian J Appl Anim Sci* 3: 423-430.
24. Silanikove N (2000) Effects of heat stress on the welfare of extensively managed domestic ruminants. *Livest Prod Sci* 67: 1-18.
25. Rasooli A, Nouri M, Khadjeh GH, Rasekh A (2004) The influences of seasonal variations on thyroid activity and some biochemical parameters of cattle. *Iran J Vet Res* 5: 1383-1391.
26. Rohn M, Tenhagen BA, Hofmann W (2004) Survival of dairy cows after surgery to correct abomasal displacement. Clinical and laboratory parameters and overall survival. *J Vet Med A* 51: 294-299.
27. Moore F (1997) Interpreting serum chemistry profiles in dairy cows. *Vet Med* 92: 903-912.
28. Adedeji O (1992) Rapid Interpretation of Routine Clinical Laboratory Tests. 1st ed Nigeria Samaru Zaria.
29. Opara M, Udevi N, Okoli I (2010) Haematological parameters and blood chemistry of apparently healthy West African Dwarf. *New York sci j* 3: 68-72.
30. Feldman B, Zink J, Jain N (2002) Schalm's Veterinary Hematology.
31. Piccione G, Messina V, Marafioti S, Casella S (2012) Changes of some haematochemical parameters in dairy cows during late gestation post-partum lactation and dry periods. *Vet Med Zoot* 58: 59-64.
32. Habeeb AA, Marai IFM, Kamal TH (1992). Heat stress. In: *Farm Animals and the Environment*.
33. Vaidya, MB, Vaghari PM, Patel BM (1970) Haematological constituents of blood of goats. *Indian Vet J* 47: 642-647.
34. Bhatta M, Das D, Ghosh PR (2013) Seasonal Variation in Erythrocytic Indices of Black Bengal Goats *Capra aegagrus hircus* in Purulia West Bengal. *Ind J Anim Health* 52: 43-48.
35. Pospisil J, Kase F, Vahala J (1987) Basic haematological values in the Cameroon goat *Capra hircus*. *Com Biochem and Physio* 88: 451-454.
36. Kaushish S, Bhattia DC, Arora KL (2013) Studies on the adaptability of sheep to subtropical climate and seasonal changes in rectal temperature cardiorespiratory and haematological attributes of Nali sheep. *Ind Vet J* 33: 760-765.
37. Piccione G, Caola G, Refinetti R (2007) Annual rhythmicity and maturation of physiological parameters in goats. *Res Vet Sci* 83: 239-243.





- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>