

Research Article

Determination of the Seasonal Yields of Total Fucose and Fucoïdan Yields in Brown Seaweeds (Order Fucales) Distributed along the Coast of Urla (izmir, Turkey)

Gamze Turan*, Semra Cirik, Hatice Tekogul, Edis Koru, Ulviye Karacalar and Safak Seyhaneyildiz

Department of Aquaculture, Ege University, Bornova, Izmir, Turkey

Abstract

The brown seaweeds have been recognized as a potential source of biologically active Fucoïdan, a mucilaginous sulfated polysaccharide consisting mainly of a polymer of α -L-fucose-4-sulfate but with persistent amounts of other sugars, mainly Xylose and Galactose and have been reported to have Anti-Cancer, Anti-Obesity and Anti-Inflammatory effects.

This study was carried out to determine the seasonal yields of total Fucose and Fucoïdan from brown seaweed species (Order Fucales) of *Cystoseira barbata*, *C. compressa* and *Sargassum vulgare* distributed naturally along the coast of Urla (Izmir, Turkey) using a colorimetric method.

Mean fucoïdan yields were $6.44 \pm 1.44\%$ (dry weight basis) for *Cystoseira barbata*, $3.84 \pm 1.14\%$ for *Sargassum vulgare* and $3.04 \pm 1.77\%$ for *C. compressa* (n=3). The highest Fucoïdan yields in these species were determined in the fall season ($P \leq 0.05$).

Keywords: Algae; Colorimetric determination; *Cystoseira*; Extraction; *Sargassum*

Introduction

In recent years, great research and commercial attention have been given to extraction and production of high-value biochemical from marine seaweed biomass [1]. The use of seaweed biomass has several

*Corresponding author: Gamze Turan, Department of Aquaculture, Ege University, Bornova, izmir, Turkey Tel: +90 2323115217; E-mail: gamze.turan@ege.edu.tr

Citation: Turan G, Cirik S, Tekogul H, Koru E, Karacalar U, et al. (2017) Determination of the Seasonal Yields of Total Fucose and Fucoïdan Yields in Brown Seaweeds (Order Fucales) Distributed along the Coast of Urla (izmir, Turkey). J Aquac Fisheries 1: 005.

Received: August 02, 2017; **Accepted:** October 26, 2017; **Published:** November 09, 2017

advantages, among them (a) available rich resources; (b) feasibility of growing fast in the open ocean; (c) higher photosynthetic efficiency than terrestrial biomass; (d) no dependence on freshwater and little dependence on temperature; and (e) low costs of harvest [2,3]. The chemistry of marine seaweeds and terrestrial biomass sugar polymers are different [1,4]. Seaweeds contain several unique and commercially important Phycocolloids, such as Alginate, Carrageenan and Agar-Agar [1,4]. Seaweeds contain other interesting and Phytochemically active molecules such as, Phycobilin pigments, HUFA and PUFA Fatty acids, Proteins with essential Amino acids, Vitamins and Mineral elements, which are compounds with potential applications in Food, Cosmetic, Pharmaceutical and Medical fields [3,5,6].

Brown Seaweed-derived polysaccharides such as Fucoïdan, Laminaran, Alginates and Mannitol have been studied due to their biological effectiveness as Anticoagulant, Antitumor, Antithrombotic, Anti-inflammatory, Contraceptive and Antiviral agent [7-10].

Fucoïdians, is a term that covers a family of Fucose-Containing Sulfated Polysaccharides (FCSPs), may constitute up to 25-30% of the seaweed dry weight, depending on the seaweed species and, to a lesser extent, on life history stage and season [11,12].

Fucoïdians participate in the building of cell walls and the support of their structure, as well as in releasing of spores and gametes from reproductive organs. Due to the sulfated groups, these polysaccharides can bind positively charged ions, such as K^+ , Na^+ , Ca^{2+} , Mg^{2+} and participate in ion exchange with the environment, which facilitates adaptation of seaweeds to water salinity fluctuations and to the toxic effects of heavy metals. Fucoïdians play a substantial role in morphogenesis of zygotes of Fucoïdan algae, as they participate in establishing the cell polarity and fixing the cell-division axis, thus determining the direction for the development of the rhizoidal and apical poles of young thalli [13].

Documented biological functionalities of these compounds are the activities against Hepatitis, Herpes and Human Acquired Immune Deficiency (AIDS) Viruses, the Anticoagulant, Anti-inflammatory Heparin and molecules that modify cell proliferation and adhesion and fertilization functions [13-20].

Despite all the above-mentioned benefits, research on the Fucoïdan content of Seaweeds in Turkey is limited.

The aim of the current study was to determine the sulfated polysaccharides (Fucoïdan) yields of seaweeds of *Cystoseira barbata*, *C. compressa* and *Sargassum vulgare* (Order Fucales) that are distributed on the Turkish coast. An experimental design was implemented to evaluate the Fucoïdan content and effect of the seasonal variation on the Fucoïdan yield among the species with specific goal to determine the best time of the year to obtain Fucoïdan from natural populations of these species.

Materials and Methods

Seaweed samples of *Cystoseira barbata*, *C. compressa* and *Sargassum vulgare* used in this study was collected monthly by hand along the coast of Urla (izmir Bay, Turkey; Figure 1). Approximately

10 m long study area where seaweed collected from sup-tidal rocky zone located between the port of Urla and Urla Island road. All three types of seaweed samples were collected from the same type of environment on the same dates. The studies species were collected from the dense beds through the year. *Cystoseira* species were identified according to Herbarium of Prof. Dr. Sukran CIRIK and personal communications with him.




Figure 1: The map of study area.

The collected seaweeds in the lab were cleaned from epiphytes and epizoa washed with distilled water, dried at (60°C), ground and stored in the freezer until use. Unless stated otherwise, chemicals and solvents used in this study were of Analytical Grade. Extraction and colorimetric determination of Fucose and Fucoïdan content of Seaweeds was estimated using the method described by [21], as described below.

Extraction

A 1g sample of seaweed material for each species of *Cystoseira barbata*, *C. compressa* and *Sargassum vulgare* for each sampling was accurately weighed into a 100 ml conical flask. 25 ml 0.2 N HCl was added and the suspension was heated at 70°C in a temperature-controlled water bath for 1 hour. The suspension was stirred (approximately 200 rpm) or shaken during the extraction. The flask was cooled in tap water and the particles were allowed to settle. The liquid was poured through a filter into a 200 ml conical flask. Subsequent extracts from the same sample were filtered into the same flask. The particles retained on the filter were transferred back into the extraction flask with 25 ml 0.02 N HCl. The flask was heated at 70°C for 1 hour, cooled in tap water and re-filtered. The combined extracts contain the acid-soluble, Fucose-containing polysaccharides used in the colorimetric procedure for Fucose determination.

Colorimetric determination of fucose

A 1 ml of sample extract was taken into a test tube placed in an ice-water bath and 4.5 ml dilute H₂SO₄ was added carefully and allowed the mixture cool for approximately 1 minute and mixed with a glass rod. The tube was placed in a boiling-water bath for exactly 10 minutes and then cooled in tap water to room temperature (20°C). 0.1 ml Cysteine solution was added into the tube and mixed with a glass rod and the tube was left at room temperature for 30 minutes.

The absorbance at 396 and 427 nm was read in this solution. The total Fucose, F, in milligrams was calculated from the following formula:

$$F = B \times 0.06 \times V$$

Where,

B is the difference between the two absorbancy readings (396 and 427 nm)

0.06 is a factor converting absorbance to amount of Fucose in milligrams per milliliter and

V is the total volume of the extract in milliliters

To obtain the approximate amount of Fucoïdan of the sample, the Fucose content was multiplied by a factor of 2 [21].

Statistical analysis

Fucose and fucoïdan yield data were analyzed using the General Linear Model (GLM) with ANOVA procedure using SPSS [22]. All data were expressed as mean ± standard deviation. Means of the fucose and means of the fucoïdan yield were compared among the species and seasons in the same species group using Tukey's Multiple comparison test in SPSS [22].

Results and Discussion

Total yield of Fucoïdan in dry weight (dw) averaged at 6.44±1.44% in *Cystoseira barbata*, 3.04±1.77% in *C. compressa* and 3.84±1.14% in *Sargassum vulgare* (Table 1). Mean Fucoïdan yield of *C. barbata* was significantly greater than that *C. compressa*, however no significant difference was found between the mean Fucoïdan yield of *C. barbata* and *S. vulgare* and *C. compressa* and *S. vulgare* (Table 1).

Seaweed Species	Mean*±SD	Minimum	Maximum
<i>Cystoseira barbata</i>	6.44±1.44	4.14	8.56
<i>Cystoseira compressa</i>	3.04±1.77	2.08	4.51
<i>Sargassum vulgare</i>	3.84±1.14	2.26	6.13

Table 1: Comparison of annual fucoïdan yield (% dry basis, with mean and Standard deviation) of *Cystoseira barbata*, *C. compressa* and *Sargassum vulgare* (N=12).

* = Means with common superscript letters are not significantly different (P≤ 0.05); (a, b): comparison between seaweed species.

The fucoïdan yields of seaweed species recorded here are similar to that the values in Chotigeat et al. [23], Larsen [21] and Usov et al. [24]. Chotigeat et al. [23] reported a fucoïdan content of 2.74±1.18% in dw in *Sargassum polycystum*, which compared favorably with *Pelvetica canaliculata*. Larsen (1978) reported a Fucoïdan content of 6-8% in dw in *Ascophyllum nodosum*, 9-11% in *Fucus* species, 5-20% in *Laminaria sp.*, and 20% in dw in *Pelvetia canaliculate*. Usov et al. (2001) reported the fucoïdan content of between 0.4 and 20.4% in dw in 25 different brown seaweed species and 7.7% in *Fucus evanescens* (Order Fucales).

Mean Fucoïdan yield in *Cystoseira barbata* was highest in spring and fall and lowest in winter (Table 2). Similarly, mean fucoïdan yield in *C. compressa* was highest in spring and fall and lowest in winter (Table 2). In *Sargassum vulgare*, Mean Fucoïdan yield was highest in fall and summer and lowest in winter (Table 2).

Species	Seasons*	Mean*±SD
<i>C. barbata</i>	Winter ^{bt}	4.79 ^b ±1.59
	Spring ^a	6.73 ^a ±1.53
	Summer ^{ab}	6.22 ^{ab} ±1.61
	Fall ^a	8.10 ^a ±1.60
<i>C. compressa</i>	Winter ^b	2.11 ^b ±1.02
	Spring ^a	3.58 ^a ±1.85
	Summer ^{ab}	2.94 ^{ab} ±1.37
	Fall ^a	3.55 ^a ±1.49
<i>S. vulgare</i>	Winter ^b	2.78 ^b ±1.21
	Spring ^{ab}	3.59 ^{ab} ±1.19
	Summer ^{ab}	3.96 ^{ab} ±1.10
	Fall ^a	5.32 ^a ±1.24

Table 2: Comparison of seasonal Fucoïdan yield (% dry basis, with mean and Standard deviation) of *Cystoseira barbata*, *C. compressa* and *Sargassum vulgare* (N=3).

*=Means with common superscript letters are not significantly different (P<0.05); (a, b): comparison between seaweed species.

Chotigeat et al. [23] have also reported a seasonal variation in Fucoïdan content in the seaweeds they examined, with the highest yields of Fucoïdan found in fall.

Addition to species and the season, the extraction method also is an important factor that has great effect on the Fucoïdan yields of the sea weeds showed by Wang and Cheng [25]. In the study of Wang and Cheng it was found that the fucoïdan content extracted using method I was 13.13% (of dry weight), 24.00% and 22.67% from *Sargassum glaucescens*, *Sargassum horneri* and *Laminaria japonica*, respectively. By contrast, the fucoïdan content extracted using method II was 4.20%, 4.80% and 4.64% from *S. glaucescens*, *S. horneri* and *L. japonica*, respectively. The sample sex-tracted using method II were lower in total sugar content but higher in sulfated content than those extracted using method I. Using method II yielded the highest fucose content in *S. Glaucescens* (20.89%) and *Hizikia fusiformis* (15.89%).

Conclusion

In this paper, the extraction, colorimetric determination of Fucose and Fucoïdan yields from *Cystoseira barbata*, *C. compressa* and *Sargassum vulgare* seaweed distributed along the coast of Urla (izmir, Turkey) has been successfully demonstrated. Results indicate that annual mean content of Fucoïdan was significantly different among the three seaweed species. For Fucoïdan extraction, *C. barbata* was better source than *C. compressa* and *S. vulgare* and the best period to harvest *C. barbata* was fall seasons (September, October and November).

Addition to the species and seasonal conditions, the extraction method has also a significant effect on the yields of Fucoïdan. Considering of sulfate content of Fucoïdians, it is supposed that the functional activity of fucoïdians between the three species will be similar because sulfate is well-known factor of Fucoïdan's biological function. So, each species should be considered a good source for Fucoïdan that has the activities against Hepatitis, Herpes and Human Immunodeficiency (AIDS) Viruses, Anticoagulant heparin inflammation, Cell proliferation and adhesion.

Acknowledgement

Authors thank to Ege University Scientific Research Projects for

providing funding for this work through grant BAP Project numbered 2004-SUF-002. The authors are also grateful to Dr. Amir NEORI for valuable information and suggestions in the preparation of the manuscript and to Prof. Dr. Sukran CIRIK for identification of seaweed species of *Cystoseira barbata*, *C. compressa* and *Sargassum vulgare*.

References

- McHugh DJ (2003) A guide to the seaweed industry. FAO Fisheries technical paper 441: 1-118.
- Ross AB, Jones JM, Kubacki ML, Bridgeman T (2008) Classification of macroalgae as fuel and its thermochemical behavior. *Bioresource Technology* 99: 6494-6504.
- Anastasakis K, Ross AB, Jones JM (2011) Pyrolysis behavior of the main carbohydrates of brown macro-algae. *Fuel* 90: 598-607.
- Percival E, McDowell RH (1967) Variations in the carbohydrate content of algae. In: Chemistry and enzymology of marine algal polysaccharides. Academic Press, London, UK.
- O'Sullivan L, Murphy B, McLoughlin P, Duggan P, Lawlor PG, et al. (2010) Prebiotics from marine macroalgae for human and animal health applications. *Mar Drugs* 8: 2038-2064.
- Lordan S, Ross RP, Stanton C (2011) Marine bioactives as functional food ingredients: potential to reduce the incidence of chronic diseases. *Mar Drugs* 9: 1056-1100.
- Bhakuni DS, Rawat DS (2005) Bioactive metabolites of marine algae, fungi and bacteria. In: Bhakuni DS, Rawat DS (eds.). *Bioactive marine natural products*. Springer, New York, USA. pg no: 1-25.
- Imbs TI, Krasovskaya NP, Ermakova SP, Makarieva TN, Shevchenko NM, et al. (2009) Comparative study of chemical composition and antitumor activity of aqueous-ethanol extracts of brown algae *Laminaria cichorioïdes*, *Costaria costata* and *Fucus evanesces*. *Russian Journal of Marine Biology* 35: 164-170.
- Wang J, Liu L, Zhang Q, Zhang Z, Qi H, et al. (2009) Synthesized oversulfated, acetylated and benzoylated derivatives of fucoïdan extracted from *Laminaria japonica* and their potential antioxidant activity *in vitro*. *Food Chemistry* 114: 1285-1290.
- Mestechkina NM, Shcherbukhin VD (2010) Sulfated polysaccharides and their anticoagulant activity: A review. *Applied Biochemistry and Microbiology* 46: 267-273.
- Ale MT, Meyer AS (2013) Fucoïdians from brown seaweeds: an update on structures, extraction techniques and use of enzymes as tools for structural elucidation. *RSC Advances* 3: 8131-8141.
- Kusaykin M, Bakunina I, Sova V, Ermakova S, Kuznetsova T, et al. (2008) Structure, biological activity and enzymatic transformation of fucoïdians from the brown seaweeds. *Biotechnol J* 3: 904-915.
- Skriptsova, AV (2015) Fucoïdians of brown algae: biosynthesis, localization and physiological role in thallus. *Russian Journal of Marine Biology* 41: 145-156.
- Elouali M, Boissonvidal C, Durand P, Josefovic J (1993) Antitumor activity of low molecular weight fucans extracted from brown seaweed *Ascophyllum nodosum*. *Anticancer Res* 13: 2011-2019.
- Berteau O, Mulloy B (2003) Sulfated fucans, fresh perspectives: structures, functions and biological properties of sulfated fucans and an overview of enzymes active toward this class of polysaccharide. *Glycobiology* 13: 29-40.
- Ponce NM, Pujol CA, Damonte EB, Flores ML, Stortz CA (2003) Fucoïdians from the brown seaweed *Adenocystis utricularis*: extraction methods, antiviral activity and structural studies. *Carbohydr Res* 338: 153-165.

17. Queiroz KC, Medeiros VP, Queiroz LS, Abreu LR, Rocha HA, et al. (2008) Inhibition of reverse transcriptase activity of HIV by polysaccharides of brown algae. *Biomed Pharmacother* 62: 303-307.
18. Foley SA, Szegezdi E, Mulloy B, Samali A, Tuohy MG (2011) An unfractionated fucoidan from *Ascophyllum nodosum*: extraction, characterization and apoptotic effects *in vitro*. *J Nat Prod* 74: 1851-1861.
19. Qu G, Liu X, Wang D, Yuan Y, Han L (2014) Isolation and characterization of fucoidans from five brown algae and evaluation of their antioxidant activity. *Journal of Ocean University of China* 5: 851-856.
20. Zayed A, Muffler K, Hahn T, Rupp S, Finkelmeier D, et al. (2016) Physicochemical and biological characterization of fucoidan from *fucus vesiculosus* purified by dye affinity chromatography. *Mar Drugs* 14.
21. Larsen B (1978) Fucoidan. In: Hellebust JA, Craigie JS (eds.). *Handbook of Phycological Methods*. University Press, Cambridge, Canada. Pg no: 151-156.
22. Ozdamar K (1997) *Package Program and Statistical Data Analysis 1*. Anatolia University Publishing, Publication number 1001, Faculty of Science Publication number 11, Eskisehir (In Turkish).
23. Chotigeat W, Tongsupa S, Supamataya K, Phongdara A (2004) Effect of fucoidan on disease resistance of black tiger shrimp. *Aquaculture* 233: 23-30.
24. Usov AI, Smirnova, GP, Klochkova, NG (2001) Polysaccharides of Algae: 551. Polysaccharide composition of several brown algae from kamchatka. *Russian Journal of Bioorganic Chemistry* 27: 395-399.
25. Wang CY, Chen YC (2016) Extraction and characterization of fucoidan from six brown macroalgae. *Journal of Marine Science and Technology*.