

# HSOA Journal of Addiction & Addictive Disorders

# **Research Article**

# Patterns of Admission for Alcohol Dependence Syndrome Treatment among Prisoners Over 18 Years in Poland

#### Katarzyna Czechowska<sup>1\*</sup>and Kamila Kujawa<sup>2</sup>

<sup>1</sup>Department of Health Service, Penitentiary Institution in Czerwony Bór, Łomża, Poland

<sup>2</sup>Ursynów Stokłosy Clinic in Warsaw, Specialist for Analysis, Warsaw, Poland

#### **Abstract**

#### **Background**

Alcohol dependency syndrome is a disease often found among persons staying in prison. Nowadays, in most cases, the patients are treated in prison therapeutic departments and in prison wards. People qualified for therapy wait for the admission in prisons and are transported to the therapy units as soon as the place is unlocked. Addicts are treated by the prison psychiatrists, psychotherapists and psychologists.

#### Methods

Demographics data from Information and Statistics Bureau CZSW were analyzed over a period of 18 years. This focused on number of admissions and type of therapy (inmates treated in prison therapy department; prisoners awaiting admission to prison therapy department; inmates treated outside the therapy department).

#### Results

Over 18 years, 29.002 statistical data were collected describing the number of prisoners diagnosed as alcohol addicts and qualified for therapy. Between 2001 and 2002 there was a statistically significant decrease in the number of prisoners treated for alcohol dependence syndrome (p <0.05). In the years 2002 - 2018 there was an increase in the number of prisoners treated due to alcohol dependence syndrome (p <0.05). In 2012, a statistically significant

\*Corresponding author: Katarzyna Czechowska, Department of Health Service, Penitentiary Institution in Czerwony Bór, Łomża, Poland, Tel: +48 501198687; E-mail: katarzyna.czechowska@gmail.com

**Citation:** Czechowska K, Kujawa K (2020) Patterns of Admission for Alcohol Dependence Syndrome Treatment among Prisoners Over 18 Years in Poland. J Addict Addictv Disord 7: 36.

Received: February 15, 2020; Accepted: February 24, 2020; Published: March 02, 2020

**Copyright:** © 2020 Czechowska K, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

decrease was observed in prisoners treated for alcoholism (p <0.05). Admissions were higher among prisoners in 2013 treated in connection with alcoholism (p <0.05). In the period 2013 - 2018, the changes in the number of prisoners treated due to alcohol dependence syndrome were not statistically significant.

#### Conclusion

There has been a significant change in admissions for treatment for alcohol dependence syndrome over the past 18 years among prisoners staying in Polish penitentiary units.

**Keywords:** Addictions; Alcohol dependence syndrome, Prisoners; Prison health care

#### Introduction

Alcohol dependency syndrome is a disorder, whose diagnosis according to the ICD-10 criterion requires the presence of at least three of the following symptoms for a certain period of time during the previous year:

- 1. Strong need or urge to drink alcohol.
- 2. Difficulties in controlling the start or end of drinking or the amount of alcohol consumed.
- Occurrence of withdrawal syndrome after stopping or reducing the dose of alcohol, manifested by the characteristic symptoms of alcohol withdrawal syndrome or taking the same (or very similar substance) to reduce the severity or avoid symptoms of withdrawal syndrome.
- 4. Symptoms of tolerance, such as increasing the dose of alcohol to obtain effects that were initially caused by lower doses (a good example is people addicted to alcohol who are able to drink the amount of alcohol that could cause loss of consciousness or death in unaccustomed people).
- 5. Increasing neglect of other pleasures or interests due to drinking alcohol.
- Increased time spent on getting alcohol, drinking alcohol or regaining balance after drinking alcohol.
- 7. Continued drinking despite obvious evidence of harmful consequences such as liver damage due to heavy drinking or depressed mood following periods of drinking large amounts of alcohol [1].

The State Agency for Solving Alcohol Problems estimates that the number of people addicted to alcohol in Poland is about 800 thousand. (2% of the population) [2]. Untreated disease can lead to complications, including cirrhosis, cachexia, polyneuropathy, gastritis and the co-occurrence of financial problems in contact with people, family life disorders. Moreover, research shows that alcohol addiction affects penality increase. People under the influence of alcohol in Poland are mostly perpetrators of violence and there is a statistically significant link between car accidents and alcohol consumption [3]. Some of these people are tried by the court for causing a criminal offense resulting in a restriction of liberty. Convicted persons are sent to prison, where they might be diagnosed alcohol dependence syndrome

and treatment might be prescribed. Serving a prison sentence where human freedom is limited means that the prisoner has no influence on the choice of the attending physician and the method of treatment, so the whole prison system has the responsibility to care for the sick.

This work was created as an analysis of the number of prisoners diagnosed per alcohol dependence syndrome and qualified for treatment in prisons in Poland in the years 2001-2018. Data was obtained from the Information and Statistics Office of the Central Management of Polish Prison available on the website www.sw.gov.pl.

# **Methods and Materials**

The following epidemiological descriptive study uses secondary sources of information in the form of aggregate reports available at the Information and Statistics Office of the Central Management of Polish Prison available on the website www.sw.gov.pl. The analysis covered a group of prisoners in Poland in prisons in the years 2001-2018.


The following variables were taken for analysis: The number of prisoners addicted to alcohol staying in wards, the number of prisoners addicted to alcohol waiting to be admitted to the ward, the number of prisoners addicted to alcohol treated outside the ward and the total number of prisoners addicted to alcohol as a percentage of the total number of prisoners.

### **Results**

#### Characteristics of the studied population

In 2001, the number of prisoners treated for alcohol dependency syndrome was 2,495 people and constituted 3.13% of the total population of prisoners, while at the end of 2018 this number was 2,127 and this year the percentage of prisoners treated for alcohol dependence syndrome in relation to the total number of prisoners was 2.95%.

The trend graph of the percentage of prisoners treated for alcohol dependence as a percentage of all prisoners is presented in figure 1.




**Figure 1:** Percentage of prisoners treated in prisons for alcohol dependence syndrome in Poland in the years 2001-2018, statistically significant decreases are marked in red; Source: Own study based on materials made available by the Information and Statistics Office of the Central Statistical Office.

In 2002 compared to 2001 there was a statistically significant decrease in the number of prisoners treated for alcoholism-a decrease from 3.13% of all prisoners to 0.67% of the total number of prisoners.

In the period 2002 - 2012 there was a constant slight increase in the percentage of prisoners treated in relation to the total number of prisoners - at the end of 2011 this percentage reached the level of 2.27%. Year 2012 presents a significant decrease in the percentage of prisoners treated in relation to the total number of prisoners - from 2.27% to 1.78% at the end of 2012 year, followed by an increase to 2.67% in 2013. In the years 2014 - 2018 changes in the percentage ratio of the number of prisoners treated for alcoholism in relation to the total number of prisoners are not statistically significant - they range between 2.81% and 2.95%.

The trend in the number of prisoners in Poland treated for alcohol dependence syndrome in absolute values in the years 2001 - 2018 is presented in figure 2.



**Figure 2:** Total prisoners addicted to alcohol treated in prisons in Poland in the years 2001 - 2018, statistically significant increases are marked in blue, statistically significant decreases in red; Source: Own study based on materials made available by the Information and Statistics Office of the Central Statistical Office.

In the analyzed period, with the exception of 2001, which is significantly different from the other analyzed years, the average number of prisoners treated for alcohol dependence was 1.559 prisoners and increased in the examined period from the number of 539 prisoners treated in 2002 to the number of 2.127 prisoners treated in 2018, which is identified as a statistically significant increase.

In 2002 compared to 2001 there was a statistically significant decrease in the number of prisoners treated for alcohol dependence syndrome-from the level of 2,495 people to the level of 539 people (almost five-fold decrease).

Starting from 2002, the total number of prisoners treated for alcohol dependence in prisons in Poland showed an upward trend except for 2008-2010 and 2013-2018, when the increases were not statistically significant.

In 2012, a statistically significant decrease was observed in prisoners treated in connection with alcohol dependence syndrome. As observed in the table 1 below, it was observed that the decrease was related to the decrease of prisoners qualified for the group awaiting admission to the department at the end of 2012 (a decrease of 416 people). The appearance in 2013 of a group of inmates treated outside the department in statistically significant numbers (671 people at the end of 2013 compared to 3 people at the end of 2012) constituted a statistically significant increase and contributed to a statistically significant increase year-over-year of the total number of prisoners treated in Polish prisons.

#### Characteristics of the studied population by subgroups

The number of prisoners treated in connection with the alcohol dependence syndrome staying in treatment departments, the number of prisoners waiting to be admitted to the department and the number of prisoners treated for the alcohol dependence syndrome outside the department in the years 2001-2018 is presented in table 1 below (statistically significant year-on-year increases are marked in blue, statistically significant year-on-year decreases are marked in red).

Date	Inmates Treated in Departments	Inmates Waiting for Admission to the Department	Inmates Treat- ed Outside the Department	Inmates Treated for Alcohol Depen- dence Syndrome - Total
31.12.2018	1108	546	473	2127
31.12.2017	1167	502	467	2136
31.12.2016	1139	439	433	2011
31.12.2015	1088	428	519	2035
31.12.2014	1008	496	564	2068
31.12.2013	1015	421	671	2107
31.12.2012	1075	421	2	1498
31.12.2011	1008	837	1	1846
31.12.2010	943	727	2	1672
31.12.2009	881	721	0	1602
31.12.2008	792	722	1	1515
31.12.2007	762	627	0	1389
31.12.2006	706	492	2	1200
31.12.2005	635	444	1	1080
31.12.2004	546	399	3	948
31.12.2003	460	259	15	734
31.12.2002	396	119	24	539
31.12.2001	384	180	1931	2495

**Table 1:** The number of prisoners treated per alcohol dependence syndrome in the years 2001-2018 broken down into subgroups; Source: Own study based on information provided by the Information and Statistics Office of the Central Statistical Office; Year-on-year increases statistically significant are marked in blue, statistically significant year-on-year decreases are marked in red.


In 2001, out of the total amount of 2,495 prisoners treated for alcohol dependence, 77% (1.931 prisoners) were treated outside the department; in the following year the number of prisoners treated outside the department decreases by 98% to the number of 24 people and this is a statistically significant decrease.

The absolute number of prisoners treated in connection with alcohol dependence syndrome increased statistically significantly in the years 2003-2008, 2011 and 2013. In 2012, the number of inmates treated for alcohol dependence syndrome fell by 348 people (23%) and this is a statistically significant decrease - the decrease in 2012 was mainly related due to almost a half of the number of prisoners awaiting admission to the department, while the number of inmates remaining in the department and inmates treated outside the department remained unchanged. In 2013, there was a statistically significant increase in the number of inmates treated outside the department (from 2 inmates to 671 inmates), which contributed to a statistically significant increase in the number of treated inmates in total. In the remaining years analyzed, the number of prisoners treated in connection with alcohol dependence syndrome remained at a similar

level-changes in numbers in the years 2014 - 2018 were not statistically significant.

The largest subgroup among prisoners treated for alcohol dependence syndrome was a subgroup of inmates located in the department and, with the exception of 2001, it was the subgroup with the highest amount of inmates treated for alcohol dependence syndrome.

The chart of the trend of the number of prisoners treated for alcohol dependence syndrome treated in the department in the years 2001 - 2018 is presented in figure 3.



**Figure 3:** Graph of the number of prisoners treated for alcohol dependence in the years 2001 - 2018; Source: Own study based on information provided by the Information and Statistics Office of the Central School of Statistics (blue indicates statistically significant increases).


In the analyzed period, the average number of prisoners treated for alcohol dependence syndrome and treated in the department was 840 people and, with exception of 2001, constituted in all years the largest subgroup among all those treated for alcohol dependence syndrome (on average 57% of all prisoners in the period 2002 - 2018). The number of inmates staying in the analyzed period in the treatment department of alcohol addicts showed a statistically significant upward trend - on average in the period under study it was an increase of 7% in the years 2002 - 2018, statistically significant increases year-to-year were recorded in the years 2003 - 2005 and in 2009.

The second analyzed subgroup of prisoners treated for alcohol dependence syndrome was a subgroup of prisoners awaiting for admission to the department. This was a subgroup significantly smaller than the first subgroup - in the years 2002 - 2018 it was on average 487 prisoners and it is on average 34% of all prisoners treated for alcohol dependence syndrome.

The chart showing the trend of the number of prisoners treated for alcohol dependence awaiting admission to the department in the years 2001 - 2018 is shown in figure 4.


The most statistically significant decrease occurred in 2002 - the number of inmates waiting for admission dropped from 180 to 119 persons and in 2012-the number of inmates waiting admission to the department dropped from 837 to 421 (almost by half), a statistically significant decrease was also recorded in 2015 (decrease from 496 to 428 prisoners).

The third, on average, least numerous subgroup of prisoners treated for alcohol dependence is the subgroup of prisoners treated outside the department.



**Figure 4:** Graph of the number of prisoners awaiting admission to the department in the years 2001 - 2018; Source: Own study based on information provided by the Information and Statistics Office of the Central School of Statistics (blue indicates statistically significant increases, red - statistically significant decreases).

The trend graph for the number of prisoners treated outside the ward for alcohol dependence syndrome in the years 2001-2018 is presented in figure 5.



**Figure 5:** Graph of the number of prisoners treated for alcohol dependence syndrome outside department in the years 2001-2018; Source: Own study based on information provided by the Information and Statistics Office of the Central School of Statistics (blue indicates statistically significant increases, red-statistically significant decreases).

At the end of 2001, the number of inmates treated outside the department was 1,931 people and was the most numerous subgroup of inmates treated (77% of all inmates treated for alcohol dependence syndrome). From 2002 to 2012, the number of inmates treated outside the department decreased significantly (24 in 2002 and 15 in 2003) and ranged from 0 to 3 inmates in the period 2004 - 2018, which constitutes statistically insignificant values; statistically significant values amounting to an average of 25% of the total number of prisoners with alcohol dependence syndrome began to be noted from 2013, in 2013 - 2014 and in 2015 - 2016 the number of prisoners treated outside the department increased statistically significantly and stood at an average level of 521 prisoners.

### **Discussion**

According to data from the Central Statistical Office of Poland in 2017, it appears that in 2016, over 4% of all Poles were treated in counseling centers for people with mental disorders, addicted to alcohol and other substances [4]. It is estimated that around 800,000 people in Poland may be addicted to alcohol, while the number of people abusing alcohol may be reaching 2 million [5]. Among inmates, it is

estimated that the problem of alcohol addiction is much larger. It is estimated that up to 30% of people who come to prison are alcohol addicts, of which the number may be even higher for penitentiary recidivists [6]. Studies conducted by the author during 10 years of work in prison showed that among prisoners about 30% of men were under 30 years old. Research on the trend in alcohol use behavior patterns among U.S. adults ages 18 - 65 showed that prevalences of the Infrequent Heavy Episodic and Extreme Drinkers peaked around ages 22 - 24 [7]. The above observations may explain the increased problem of alcoholism among prisoners. Education and the environment in which a person lives can have an impact on his future. Lack of interest of parents / guardians in a child, exposure to aggression, lack of resourcefulness of parents / guardians and the presence of alcohol from an early age can affect both subsequent alcohol abuse leading to addiction, as well as the commission of crimes resulting in a penalty of imprisonment [8]. The study conducted by The National Monitoring of Adolescent Prescription Stimulants Study showed that 6.8% of youth within a month used prescription stimulants. Part of this group admitted that they used these substances for non-medical purposes and were in close relationships with people who endorse binge drinking [9]. In addition to alcohol dependence, the prison population is characterized by the problem of using other stimulants, including injection drugs. The use of unsterilized equipment and a lack of knowledge about cross-infection results in an increased risk of developing hepatitis C [10]. Studies in Northern California have shown that people in the "increasing group of marijuana use" consumed alcohol more frequently [11]. The Alcohol Use Disorders Identification Test (AUDIT) performed among prisoners identifying alcohol consumption disorders is a useful tool for detecting alcohol use among prisoners and at the same time it can identify other drug problems [12]. Alcohol consumption plays an important role in aggressive behavior and is recognized as a driving force for committing crimes [13]. In 2018, the report 'Towards national measures of alcohol-related crime' was created, in which an important role in heavy crime is attributed to the component of excessive alcohol consumption as the main driving force of the phenomenon [14]. The high number of people addicted to alcohol imposes an obligation to the Polish prison system for the development of effective treatment methods. Over the years, national legislation has changed with intention to allow prisoners to qualify for therapy - the goal of the changes ways to allow as many prisoners as possible to receive individualized forms of therapy. Undoubtedly, in the conditions of prison isolation it is easier to maintain alcohol abstinence than outside the prison isolation and therapeutic programs provided to inmates in therapeutic departments practically do not differ from those carried out outside the prison isolation [15].

There are 33 therapeutic wards for alcohol-dependent prisoners in Polish prisons and detention centers, which have 1,187 places in total [16]. According to data from the Central Board of the Prison Service in Poland, alcohol addicts are serving a prison sentence in, among others, therapeutic departments of prisons in Zamość, Wronki, Radom, Opole Lubelskie [17]. Therapy programs for prisoners with alcohol dependence syndrome usually last about 3 months, during which patients under appropriate conditions undergo targeted therapy by qualified personnel [18]. It is worth noting that over the years, along with the increase in demand, the number of such centers increases and the largest group of inmates qualified for therapy are people addicted to alcohol [19]. Diagnosis of diseases during imprisonment is made by prison psychiatry specialists and referral to therapy is made by psychiatrists or psychologists after doctor's consultation.

Alcohol dependence syndrome in Polish prisons is diagnosed on the basis of diagnostic criteria contained in the Classification of Psychiatric Disorders and Behavioral Disorders ICD-10 and screening tests, including the diagnostic questionnaire for alcohol addiction syndrome according to Dr. Woronowicz [20,21].

Before the prisoner attends a specialist consultation to a psychiatrist or goes under the care of the prison psychologist, he undergoes a routine medical examination during the first three days of his stay in prison. Only after this visit can he be assigned to the target residential cell. During the first examination, the prison doctor determines whether the prisoner may be addicted to alcohol or whether he is at risk of having a post withdrawal syndrome during his stay due to abrupt withdrawal of alcohol. Post withdrawal syndrome may endanger the patient's life. Hyperactivity, hypotension and compensatory tachycardia are one of the possible complications that can be successfully treated with appropriate medications [22]. In Polish prison, a person deprived of liberty is supposed to be provided with medical treatment from the beginning of the illness until healing.

The study analyzed data on the treatment of people diagnosed with alcohol dependence syndrome in prison from 2001 to 2018. In 2001, the number of prisoners treated for alcohol dependence syndrome was 2,495 people and constituted 3.13% of the entire population of prisoners. In 2002, the number of inmates treated for alcohol dependence syndrome dropped dramatically to the value of 539 (0.67% of all prisoners) and in the following years it gradually increased until 2018, in which a number of 2,127 was reached, and the percentage of prisoners treated for alcohol dependence in relation to the total number of prisoners was 2.95%.

Between 2001 and 2002 there was a nearly 5 - fold decrease in the number of prisoners treated for alcohol dependence syndrome from 2.495 to 539 (from the percentage of 3.13% of all prisoners to 0.67% of all prisoners respectively). This decrease occurred due to a statistically significant decrease in the number of prisoners treated outside the ward-a total decrease of 1,956 prisoners, including a decrease in the number of prisoners treated outside the ward by 1,907 people (decrease by 98% in the respective sub - group). This change was associated with the entry into force of the Regulation of the Minister of Justice on ways of penitentiary interactions in prisons and detention centers [23]. The new legal document assumed that the vast majority of people qualified for the treatment of alcohol dependence syndrome was directed to have therapy in therapeutic departments. As a consequence of this decision, a gradual increase in the number of people waiting for admission to the department in the period 2002 - 2011 was noted. In 2002, 119 prisoners were awaiting transport to the therapeutic department, while in 2011 the number increased to 837 prisoners. As a result, a similar increase in the total number of inmates treated for alcohol dependence syndrome treated in the department was observed. In the years 2002 - 2018, the average number of prisoners treated for alcohol dependence was 1.559 prisoners and increased in the period under review from the number of 539 prisoners treated in 2002 to the number of 2.127 prisoners treated in 2018, which was a statistically significant increase (respectively, 0.67% of all inmates at the end of 2002 and 2.95% at the end of 2018, almost fourfold increase).

In 2012, there was a change in the regulation on conducting penitentiary interactions in prisons and detention centers. According to

the new guidelines, the therapeutic team will implement specialist impacts for convicts qualified to the therapeutic system in the therapeutic department. The possibility of intervention outside the therapeutic department by specialized staff has also been introduced. While an inmate qualified to serve a prison sentence in a therapeutic system may be subject to individualized interactions according to a developed individual therapeutic program, group therapeutic interactions in the therapeutic system outside the therapeutic department may additionally be conducted. The possibility of conducting treatment outside the therapeutic department is diagnosed and recommended by the psychologist based on the doctor's opinion or based on the psychopenitentiary judgment [24].

The above decisions had an impact on the results of the statistical analysis of the Central Board of Prison Services. According to data, in 2012 a statistically significant decrease was observed in prisoners treated for alcoholism - the percentage of prisoners treated for alcohol dependence fell from 2.27% to 1.78%. The decrease in 2012 was associated with a decrease of 416 people / 49%, the number of prisoners qualified for the group awaiting admission to the department. Then, in 2013 there was a significant increase in the number of prisoners treated for alcohol dependence (an increase of 609 people, from 1.78% of all prisoners to 2.67% of all prisoners) and this was due to the emergence of a subgroup of prisoners treated outside the department, which was a statistically significant number (671 people at the end of 2013 compared to 3 people at the end of 2012, which is a statistically significant increase). In the following years from 2014 to 2018, changes in the percentage ratio of the number of prisoners treated for alcoholism in relation to the total number of prisoners were not statistically significant - they ranged between 2.81% and 2.95%.

Based on the analyzed data, it can be seen that alcohol addiction is a serious health problem among the population of Polish prisoners. Considering the numbers given earlier, despite the fact that the number of prisoners diagnosed with alcohol dependence syndrome is many times higher than that in the general population of the country, still not many people in prison are undergoing therapy. At the same time, the fact that there is a problem of the growing number of people referred for therapy but waiting for the vacancy suggests that there is a lack of staffing or accommodation in the Polish prison system affecting the ever - increasing number of patients in need of therapy. This is not an isolated problem only for the Polish Prison. Research conducted in Australia and Ireland confirms both the greater number of people addicted to alcohol among the prisoners compared to the general population, and the need to expand the health care offer in response to the need for personalized treatment [25,26].

Persons deprived of freedom belong to a group that requires careful analysis in terms of health status, because they often come from dysfunctional environments [27]. Similar data applies to the studied groups with alcohol consumption disorders. Lower education, lower income and lower socioeconomic status of the environment were associated with an increased risk of these disorders [28]. In Poland, there are very few publications investigating the occurrence of chronic diseases among prisoners, which include alcohol dependence syndrome, while those that arose show that the health status of the prison population is unsatisfactory.

# **Summary**

The resulting work is the first study analyzing the ways of conducting therapy among persons deprived of their liberty over the

years using statistical data from the Central Office of the Prison Service Boards carried out in Poland. For this reason, it is advisable to conduct further epidemiological studies, the result of which will be to determine the main health and organizational problems, which will translate into the health policy of the Polish prison.

# Strengths and Limitations of This Study

- The use of 18 year statistical documentation in the study allowed access to a database of prisoners who were in prison and were qualified for treatment.
- The nature of the data is susceptible to incomplete or incorrect medical records and coding.

# **Declaration of Conflicting Interests**

The authors declare that there is no conflict of interest.

# **Ethical Approval**

The descriptive study did not require the approval of the bioethics committee.

# **Funding**

This research received no specific grant from any funding agency in the public, commercial, or not - for - profit sectors.

### References

- Iwanicka KA, Olajossy M (2015) The concept of alcohol craving. Psychiatr Pol 49: 295-304.
- Woronowicz B (2017) Kliniczne i społeczne aspekty uzależnienia od alkoholu. Wyzwania 4 indd 5: 165-188.
- 3. Motyka M, Frackowiak M (2015) Characteristics of social problems associated with alcohol abuse. Hygeia Public Health 50: 314-322.
- Zyra M, Malesa E (2018) Health and Healthcare in 2017, Statistics Poland, Social Surveys Department, Statistical Office in Kraków, Warsaw, Poland.
- Sowa B (2018) Alcohol consumption in Poland and its socioeconomic effects in 2001-2016. Rynek-Społeczeństwo-Kultura 29: 29-33.
- Zajęcka B (2008) Therapy of people addicted to alcohol in penitentiary units. Prace Akademii im Jana Długosza w Częstochowie Seria Pedagogika 17: 125-140.
- Bray BC, Lanza CT, Dziak JJ (2019) Age trends in alcohol use behavior patterns among U.S. adults ages 18-65. Drug Alcohol Depend 205: 107689.
- Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, et al. (1998) Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) study. Am J Prev Med 14: 245-258.
- Wang Y, Cottler LB, Striley CW (2015) Differentiating patterns of prescription stimulant medical and nonmedical use among youth 10-18 years of age. Drug Alcohol Depend 157: 83-89.
- Kinner SA, Jenkinson R et al. (2012) High-risk drug-use practices among a large sample of Australian prisoners. Drug Alcohol Depend 126: 156-160.

- Campbell CI, Sterling S, Chi FW, Kline-Simon AH (2016) Marijuana use and service utilization among adolescents 7 years post substance use treatment. Drug alcohol Depend 168: 1-7.
- Webster MJ, Staton-Tindall M, Oser C, Lukefeld C, Duvall J, et al. (2014) Brief screening for alcohol and drug problems among prisoners. Drug and Alcohol Dependence 140: 239.
- Snowden AJ (2019) Exploring violence: The role of neighborhood characteristics, alcohol outlets, and other micro-places. Soc Sci Res 82: 181-194
- Morgan A, Lyneham S, Davy D, Coughlan M (2019) AIC reports Research Report 12 Towards national measures of alcohol-related crime. Australian Institute of Criminology 2018 ISSN (Online) 2206-7280.
- Żywucka-Kozłowska E (2017) Prison sentence in the Therapeutic System. Kortowski Przegląd Prawniczy 4: 78-83.
- Krakowska E (2020) Interactions with prisoners with an alcohol problem. Central Board of the Prison Service, Warsaw, Poland.
- Structure of the Prison Service in Poland. Central Board of the Prison Service, Warsaw, Poland.
- Kudlak G (2016) Personality factors and the effectiveness of addiction therapy in prison for three personality types. Profilaktyka Społeczna i Resocjalizacja 29: 7-34.
- Majcherczyk A (2007) Peculiar features of prison-based treatment programmes for inmates addicted to psychoactive substances. Alkoholizm i Narkomania 20: 321-342.
- The ICD-10 Classification of Mental and Behavioural Disorders. Clinical description and diagnostic guidelines. World Health Organization, Geneva, Switzerland.
- Frackowiak M, Motyka M (2015) Alcohol dependence syndrome: Characteristics, development phases, methods of diagnosis. Probl Hig Epidemiol 96: 315-320.
- Aljerf L, AlMasri N (2018) Beyond pain, fear, withdrawal-findings, and problems involving change - treatment and application for a chronic addiction on alcohol do not end. Drug Designing & Intellectual Properties International Journal 2: 1-9.
- ISAP (2002) Regulation of the Minister of Justice of April 16, 2002 on ways of conducting penitentiary interactions in prisons and detention centers. Journal of Laws 58: 537.
- 24. ISAP (2012) Ordinance of the Minister of Justice of November 26, 2012 amending the ordinance on ways of conducting penitentiary interactions in prisons and detention centers. Journal of Laws 1409.
- Cannon A, Nally F, Collins A, Fay R, Lyons S (2019) Trends in addiction treatment in Irish prisons using national surveillance data, 2009-2014. Int J Prison Health 15: 105-113.
- Dolan K, Rodas A, Bode A (2015) Drug and alcohol use and treatment for Australian Indigenous and non-Indigenous prisoners: Demand reduction strategies. Int J Prison Health 11: 30-8
- Udo T (2019) Chronic medical conditions in U.S. adults with incarceration history. Health Psychol 38: 217-225.
- Calling S, Ohlsson H, Sundquist J, Sundquist K, Kendler KS (2019) Socioeconomic status and alcohol use disorders across the lifespan: A co-relative control study. PLoS One 14: 0224127



Advances In Industrial Biotechnology | ISSN: 2639-5665

Advances In Microbiology Research | ISSN: 2689-694X

Archives Of Surgery And Surgical Education | ISSN: 2689-3126

Archives Of Urology

Archives Of Zoological Studies | ISSN: 2640-7779

Current Trends Medical And Biological Engineering

International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X

Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276

Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292

Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370

Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594

Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X

Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562

Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608

Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879

Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397

Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751

Journal Of Aquaculture & Fisheries | ISSN: 2576-5523

Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780

Journal Of Biotech Research & Biochemistry

Journal Of Brain & Neuroscience Research

Journal Of Cancer Biology & Treatment | ISSN: 2470-7546

Journal Of Cardiology Study & Research | ISSN: 2640-768X

Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943

Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771

Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844

Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801

Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978

Journal Of Cytology & Tissue Biology | ISSN: 2378-9107

Journal Of Dairy Research & Technology | ISSN: 2688-9315

Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783

Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X

Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798

Journal Of Environmental Science Current Research | ISSN: 2643-5020

Journal Of Food Science & Nutrition | ISSN: 2470-1076

Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X

Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566

Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485

Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662

Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999

Journal Of Hospice & Palliative Medical Care

Journal Of Human Endocrinology | ISSN: 2572-9640

Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654

Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493

Journal Of Light & Laser Current Trends

Journal Of Medicine Study & Research | ISSN: 2639-5657

Journal Of Modern Chemical Sciences

Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044

Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X

Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313

Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400

Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419

Journal Of Obesity & Weight Loss | ISSN: 2473-7372

Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887

Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052

Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X

Journal Of Pathology Clinical & Medical Research

Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649

Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670

Journal Of Plant Science Current Research | ISSN: 2639-3743

Journal Of Practical & Professional Nursing | ISSN: 2639-5681

Journal Of Protein Research & Bioinformatics

Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150

Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177

Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574

Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060

Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284

Journal Of Toxicology Current Research | ISSN: 2639-3735

Journal Of Translational Science And Research

Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193

Journal Of Virology & Antivirals

Sports Medicine And Injury Care Journal | ISSN: 2689-8829

Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: https://www.heraldopenaccess.us/submit-manuscript