



Case Report

Recurrent Pregnancy Loss in Two Patients with High Density of NK Cells CD56+ and CD16+ in Ovulatory Cervical Mucus and in Endometrium (Case Report)

Zdenka Ulcova-Gallova^{1*}, Mukensnabl P² and Losan P¹

¹Genetics - Pilsen, part of Next Clinics, and Department of Gynecology and Obstetrics, Faculty of Medicine, Charles University, Pilsen, Czech Republic

²Department of Pathology, Faculty of Medicine, Charles University, Pilsen, Czech Republic

Abstract

We describe two patients who suffered from multiple repeated pregnancy losses and with normal hormonal status, humoral autoimmune responses to the panel of phospholipids, and normal karyotyping, without pathological trombophilia. Patients were without any clinical and laboratory symptoms of vaginitis at the time of ovulatory cervical mucus (OCM) sampling and endometrium study. Blood NK cells were studied by flow cytometry, uterine NK cells CD56+ , CD16+ and NK cells in OCM were identified by immunocytochemistry, aPIs by ELISA, indirect immunofluorescent method for detection of serum and OCM IgM, IgG antibodies against HHV-6 levels at the time of ovulation. All findings were normal except for high density of NK cells in OCM and in endometrium. Intramuscular immunoglobulins in special scheme was used for the treatment. Both patients became successfully pregnant, and delivered healthy babies.

Keywords: Missed abortion; NK cells; Endometrium; Ovulatory cervical mucus; Immunoglobulins

*Corresponding author: Zdenka Ulcova-Gallova, Genetics - Pilsen, part of Next Clinics, and Department of Gynecology and Obstetrics, Faculty of Medicine, Charles University, Pilsen, Czech Republic, Tel: +420 377241529; E-mail: ulcova-gallova@email.cz

Citation: Ulcova-Gallova Z, Mukensnabl P, Losan P (2019) Recurrent Pregnancy Loss in Two Patients with High Density of NK Cells CD56+ and CD16+ in Ovulatory Cervical Mucus and in Endometrium (Case Report). J Clin Immunol Immunother 5: 013.

Received: August 16, 2019; **Accepted:** August 27, 2019; **Published:** September 04, 2019

Copyright: © 2019 Ulcova-Gallova Z, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Recurrent pregnancy loss (RPL) is associated with chromosomal, hormonal, uterine abnormalities, immunological disorders but also with infectious agents, and various environmental factors that could play a serious problem in fertility failure. Autoimmune dysfunction representing antiphospholipid antibodies or natural killer cells dysbalance between mother and early embryo are involved in the pathogenesis of recurrent RPL too.

NK cells represent 15% of peripheral blood lymphocytes. The uterine natural killer cells (uNK) play the major immune role during implantation. The major leucocyte concentration is during the first trimester of pregnancy. The NK cells are localized to the place of embryo implantation. Decidual NK CD56+ produce the cytokines involved in trophoblast invasion, growth and angiogenesis. However, the altered numbers of endometrial NK cells lead to pregnancy pathogenesis such as pregnancy loss owing to the increased of embryocytotoxic cytokines reducing angiogenesis, as well as trophoblast and vascular growths [1-3].

We refer two infertile women with 4-6 missed abortions with only high density of NK cells in ovulatory cervical mucus (OCM) and endometrial tissue. The investigation of NK cells in OCM have never been studied before. We begin with our detailed diagnostic possibilities, continue with the treatment, and with the final results.

Methods

Routine examination of our two infertile women confirmed normal anatomy of their uterus, regular menstrual cycle and ovulation. Any evidence of vaginal and cervical cultures for mycoplasma, chlamydia, trichomoniasis, yeast, and HHV-6 infection was not found. Serum antisperm antibodies, thyroid, anti-zona pellucida, and panel of antiphospholipid antibodies were also negative. We also excluded clinical and laboratory symptoms of vaginitis, antiphospholipid syndrome, and congenital thrombophilic factors. We focused on their anamnestic basic data (e.g. nicotine, the length of their hormonal contraception before decision to conceive, stress). Stress factor can not be completely eliminated. Their male partners are healthy men with normal spermograms. Our two patients were completely healthy at the time of OCM sampling, and endometrium examination.

Our two referred women, ZB and SP gave a written informed consent before publication of their cases. They had normal findings of blood NK cells investigated by flow cytometry [4,5]. We focused on the density of NK cells CD56+ and CD16+ in OCM (never been studied before), and then in secretory endometrium. We also took blood and OCM samples for immunofluorescent identification of the IgM, IgG antibodies against HHV-6 to exclude viral infection.

Collection of OCM

The ovulatory phase is routinely established by pelvic ultrasonography

The conditions for OCM taking in days 12-14 of menstrual cycle are: Protective intercourse by condom to avoid semen contamination

from the last day of menstruation to patient's examination. During routine gynecological examination without lubricant, the ectocervix is gently wiped and cleaned of vaginal debris with a sterile swab. Endocervical mucus is carefully aspirated with a special syringe (Fortuna Optima, Denmark) directly into sterile glass capillaries, immediately spread on to glass slides and fixed for the next examination. Blood collection was performed at the time of ovulation, both sera were screened for the aPLs (classical ELISA).

Antibodies against HHV-6

The indirect immunofluorescent method was used for identification of IgM or IgG HHV-6 infection in serum, and in OCM by IF Viditest (fy Vidia) using FITS conjugat anti IgG and FITS conjugat anti IgM. Antibodies IgG are evaluated in titre 1:10 and 1:40, for IgM in 1:10.

The ELISA method was used for detection of sperm, zona pellucida, panel of antiphospholipid antibodies.

Examination of NK cells in endometrium and OCM

The endometrium was obtained by hysteroscopy performed on the 22nd-24th day of the menstrual cycle. We use formol fixed paraffin embedded biopsies to do a conventional and routine anti-CD56+ and anti CD16+ staining (Human CD56, Anti Human CD16, DAKO, Leica Microsystems GmbH, UK) for determination of lymphocytes CD56+ and CD16+. Then we use a microscope (Olympus, magnification 10x20) which allows us to count and calculate cells per mm² surface for OCM and endometrial tissue. We consider more than 20% of positively labelled cells into OCM, and more than 50% of labelled cells in endometrium to be positive, such as pathological indicator.

Treatment with intramuscular immunoglobulins

We developed a special scheme of immunoglobulin treatment for patients with pathological findings of NK cells in endometrium (Scheme 1). During this treatment, patients take low-molecular-weight-heparins (the duration of this treatment depends on the current hematological outcomes). We also add gestagens, and vitamins group B.

Patients

ZB

A 36-year-old patient (ZB), gravida IV, para 0 was sent to the Division of Infertility and Immunology of Reproduction at Genetics Institute, Pilsen in 2017 because of her recurrent pregnancy losses. The patient's childhood development was normal, with common pediatric infectious disease. She works as a teacher, she does not smoke. She has a slight pollen allergy. She used 15 years hormonal contraception (HC) before conception. ZB has been pregnant four times (four RPL). Anomalies of her uterus was excluded.

Laboratory findings showed the absence of antibody to sperm, and zona pellucida, to panel of eighteen various phospholipids. The levels of ANA, C3, C4, ANCA were normal. Thyroid and celiac disease, genetic factors (factor V 1691 G ≥ A (Leiden mutation), FII 20210 G ≥ A mutation, and MTHFR 1298 A ≥ C variants) were not present. OCM was normal, with the absence of sperm and HHV-6 antibodies.

Her husband displayed a normal spermogram with no sperm antibodies, and with normal intra-acrosomal enzymes examined by monoclonal antibodies. No chromosomal abnormalities were

detected in either partner. Both partners have normal karyotypes: 46, XX and 46, XY.

Density of NK CD 56+ and CD 16+ was high in OCM, and in endometrium.

Due to the high density of local NK cells, ZB was treated by intramuscular immunoglobulins according to the scheme (Table 1), and the combined treatment of LWMH, folic acid, pyridoxine, and gestagens immediately after positivity of HCG of her fifth pregnancy which was all the time normal.

| | ZB | SP |
|---------------------------|----------------------------|----------------------------|
| Age (years) | 36 | 34 |
| Menstrual cycle | regular | regular |
| Number of missed abortion | 4 | 6 |
| Stress | none | sometimes |
| Allergy | polen | none |
| HC (years) | 15 | 5 |
| Genetics examination | normal | normal |
| Vaginal infection | none | none |
| NK cells-blood | normal number | normal number |
| NK cells-OCM | high density CD56+ | high density CD56+ |
| | lower density CD16+ | lower density CD16+ |
| NK cells-endometrium | high density CD 56+ | high density CD56+ |
| | lower density CD16+ | high density CD16+ |
| Treatment | IMIG, gestagens, vitamin B | IMIG, gestagens, vitamin B |
| Delivery | son, 3050g/52 cm | daughter, 3280g/50 cm |

Table 1: Summarises all data about our two selected patients.

Abbreviations: RPL-repeated pregnancy loss, NK-nautral killer cells, HC hormonal contraception, IMIG-intramuscular immunoglobulin.

Result

She gave a spontaneous birth to her son (3050 g/52 cm).

SP

A 34-year-old patient (SP), gravida VI, para 0 was also sent to the Division of Infertility and Immunology of Reproduction at Genetics Institute, Pilsen in 2018/1 because of her recurrent pregnancy losses. Her family history and childhood were uneventful. She is secretary, with secondary education, with regular menstrual cycle, her length of HC was 5 years, non-smoker. Her genetics, hematological, and immunological profiles were completely normal. The levels of ANA, ANCA and C3 and C4, were normal.

Blood, OCM and secretory endometrium were examined on the density of NK cells CD56+, and CD16+. We found normal NK in the blood, however, the cervical and endometrial density was high, primarily NK cells CD 56+. When she successfully conceived, she was also treated by low doses of gestagens, vitamins B, LMWH, and intramuscular immunoglobulins.

She gave a birth spontaneously her daughter (3280/50) at week 40.

Discussion

Recurrent pregnancy loss effects up to 5% of couples. RPL includes genetic, endocrinological, immunological, anatomical, and environmental causes. We start with an extensive history, gynecological, and immunological examination to identify the cause of pregnancy failure.

Recent studies [1-10] show an immunological association between mother and her foetus in a normal and pathological pregnancy, where the mother is exposed to a great number of autoimmune and alloimmune antigens which modulate her immune activities.

We examined NK cells CD56+ and CD16+ together in endometrium and OCM in patients with ovulation, and with RPL [11], but until now, we have found pathological density of NK cells only in two of them, excluding the influence of HHV-6 infection.

Previously we investigated [4] association between density of NK cells CD56+ and NK cells CD16+ in the blood and secretory endometrium in patients with unexplained recurrent miscarriage. We monitored their blood NK cells by flow cytometry and NK cells in endometrium by immunohistochemistry. The cellular immunity was focused on the identification of lymphocytes CD56+ and CD16+. NK cells CD56+ and NK cells CD16+ in the blood were in normal number in contrast with high density of above mentioned cells in endometrium. Only two referred patients had present NK cells in OCM, and in endometrium together, we excluded influence of HHV-6 infection.

We know that NK cells in the blood, decidua, and also in secretory endometrium are among the most important cells of our immune defense system [1,6,8,9]. Generally, NK cells are basic heterogenous representatives of cellular immunity, belonging to the lymphocyte family. They are characterized by natural cytotoxicity participating in anti-infectious immunity, producing cytotoxic cytokines or/- are activated by these cytokines. NK cells also participate in the destruction of tumorous tissue. In the immunology of reproduction, those cells carrying CD56+ and CD16+ are very important. The extent of cytotoxicity depends on the expression of these signs. NK cells have also a regulatory function, T-regulatory lymphocytes and dendritic cells are increased at the time of implantation and, their cytotoxicity can be increased in case of Th1 prevalence. According to the density of NK cells we could determine a high or small degree of their possible cytotoxicity. NK cells have numerous characteristics. We have selected those expressing CD56+ and CD16+, very important in reproduction. We detect them routinely in endometrium and in OCM.

Repeated miscarriage has multifactorial causes. We reported incidence of high density of NK cells in endometrium and OCM in our two infertile women. Immuno-histological findings comprise increased density of CD56+ and CD16+. We demonstrated that application of lower dosages of immunoglobulin treatment was in both cases successful. Detailed diagnosis improves the chances for a live birth.

It has been shown [1-3,5-7,10] that the imbalance in local cellular immunity can lead to an impairment of immunological tolerance, to an increase of phagocytic activity, and Th1/Th17 rearrangement with increasing levels of embryocytotoxic cytokines, particularly of TNF-alpha and INF-gamma especially in women with repeated implant failure or early fetal loss.

If the dysregulation between NK cells and T-lymphocytes in endometrium is present, a cytotoxic environment occurs directly in the uterus, and this condition leads to impairment of proliferation and differentiation of the trophoblast, angiogenesis, and increased oxidative stress due to local ischemic changes. Repeated abnormalities of T-cells (increasing Th1 and Th17 responses and decreased Th2) are frequent in patients with pregnancy loss [10], who easily become

pregnant, but their pregnancy terminates with RPL. We did not find a correlation between local and peripheral NK cells [4].

Little is known about NK cells CD56+ and CD16+ in OCM. High density of NK cells CD56+ and/or CD16+ in cervicalovulatory mucus and in endometrium in only two infertile women with RPL was found. It seems, that NK cells in OCM are very rare, but it is not clear if NK cells in OCM are fixed among secretory cells of the cervical glands and cervical crypts of the uterine cervix, or whether these cells migrate from endometrium in the OCM. Evidence of NK cells in the OCM suggests that the uterine cervix belongs to the active immunological organs of the female genital tract involved in the local immune responses. The NK cells findings in this localization are very interesting for the future research. However, according to immunohistological examinations, NK cells also are present in cervical area but the question remains: are cervical NK cells fixed or migrating from the endometrium? Or, are cervical NK cells involved in latent infection such as HHV-6 [10]? which was not proven in this work.

Conclusion

We presented two cases who suffered from RPL due to underlying local autoimmune cell reaction. Treatment of both patients was mainly based on immunomodulation (immunoglobulins), and anti-coagulation (LMWH), and has led to successful births of healthy babies.

Scheme 1: Scheme of the treatment with intramuscular immunoglobulins (IMIG)

Igamplia (5ml, 800mg) divided in two doses is administered intramuscularly to two different muscle sites (data, and medical doctor's signature)

On the day of positive HCG

6th week of pregnancy

10th week of pregnancy

12th week of pregnancy

16th week of pregnancy

22th week of pregnancy

28th week of pregnancy

32th week of pregnancy-it is usually the last application of the Igamplia

After application of IMIG, our patients remain in the waiting room at least an hour and are monitored by medical personnel because of possible side effects.

References

1. Quenby S, Bates M, Doig T, Brewster J, Lewis-Jones DI, et al. (1999) Pre-implantation endometrial leucocytes in women with recurrent miscarriage. Hum Reprod 14: 2386-2391.
2. Saito S, Shima T, Inada K, Nakashima A (2013) Which types of regulatory T cells play important roles in implantation and pregnancy maintenance? Am J Reprod Immunol 69: 340-345.
3. Kuon RJ, Vomstein K, Weber M, Muller F, Seitz C, et al. (2017) The "killer cell story" in recurrent miscarriage: Association between activated peripheral lymphocytes and uterine natural killer cells. J Reprod Immunol 119: 9-14.

4. Ulcova-Gallova Z, Pesek M, Chaloupka P, Mukensnabl P, Haschova M, et al. (2017) Screening of endometrial NK cells in selected infertile patients. First part-methods and current results. *Ceska Gynekol* 82: 366-371.
5. Kwak-Kim J, Bao S, Lee SK, Kim JW, Gilman-Sachs A (2014) Immunological modes of pregnancy loss: Inflammation, immune effectors, and stress. *Am J Reprod Immunol* 72: 129-140.
6. Tang AW, Alfrevic Z, Quenby S (2011) Natural killer cells and pregnancy outcomes in women with recurrent miscarriages and infertility: A systemic review. *Human Reprod* 26: 1971-1980.
7. Vassiliadou N, Searle RF, Bulmer JN (1999) Elevated expression of activation molecules by decidual lymphocytes in women suffering spontaneous early pregnancy loss. *Hum Reprod* 14: 1194-2000.
8. Lee SK, Na BJ, Kim JY, Hur SE, Lee M, et al. (2013) Determination of clinical cellular immune markers in women with recurrent pregnancy loss. *Am J Reprod Immunol* 70: 398-411.
9. Seshadri S, Sunkara SK (2014) Natural killer cells in female infertility and recurrent miscarriage: A systematic review and metaanalysis. *Human Reprod Update* 20: 429-438.
10. Coulam CB, Bilal M, Salazar Garcia MD, Katukurundage D, Elazzamz H, et al. (2018) Prevalence of HHV-6 in endometrium from women with recurrent implantation failure. *Am J Reprod Immunol* 80: e12862.
11. Ulcova-Gallova Z, Mukensnabl P, Haschova M, Pesek M, Chaloupka P, et al. (2019) NK cells not only in endometrium but also in ovulatory cervical mucus in patients with decreased fertility. *Ceska Gynekol* 34: 184-189.



- Journal of Anesthesia & Clinical Care
Journal of Addiction & Addictive Disorders
Advances in Microbiology Research
Advances in Industrial Biotechnology
Journal of Agronomy & Agricultural Science
Journal of AIDS Clinical Research & STDs
Journal of Alcoholism, Drug Abuse & Substance Dependence
Journal of Allergy Disorders & Therapy
Journal of Alternative, Complementary & Integrative Medicine
Journal of Alzheimer's & Neurodegenerative Diseases
Journal of Angiology & Vascular Surgery
Journal of Animal Research & Veterinary Science
Archives of Zoological Studies
Archives of Urology
Journal of Atmospheric & Earth-Sciences
Journal of Aquaculture & Fisheries
Journal of Biotech Research & Biochemistry
Journal of Brain & Neuroscience Research
Journal of Cancer Biology & Treatment
Journal of Cardiology: Study & Research
Journal of Cell Biology & Cell Metabolism
Journal of Clinical Dermatology & Therapy
Journal of Clinical Immunology & Immunotherapy
Journal of Clinical Studies & Medical Case Reports
Journal of Community Medicine & Public Health Care
Current Trends: Medical & Biological Engineering
Journal of Cytology & Tissue Biology
Journal of Dentistry: Oral Health & Cosmesis
Journal of Diabetes & Metabolic Disorders
Journal of Dairy Research & Technology
Journal of Emergency Medicine Trauma & Surgical Care
Journal of Environmental Science: Current Research
Journal of Food Science & Nutrition
Journal of Forensic, Legal & Investigative Sciences
Journal of Gastroenterology & Hepatology Research
Journal of Gerontology & Geriatric Medicine
Journal of Genetics & Genomic Sciences
Journal of Hematology, Blood Transfusion & Disorders
Journal of Human Endocrinology
Journal of Hospice & Palliative Medical Care
Journal of Internal Medicine & Primary Healthcare
Journal of Infectious & Non Infectious Diseases
Journal of Light & Laser: Current Trends
Journal of Modern Chemical Sciences
Journal of Medicine: Study & Research
Journal of Nanotechnology: Nanomedicine & Nanobiotechnology
Journal of Neonatology & Clinical Pediatrics
Journal of Nephrology & Renal Therapy
Journal of Non Invasive Vascular Investigation
Journal of Nuclear Medicine, Radiology & Radiation Therapy
Journal of Obesity & Weight Loss
Journal of Orthopedic Research & Physiotherapy
Journal of Otolaryngology, Head & Neck Surgery
Journal of Protein Research & Bioinformatics
Journal of Pathology Clinical & Medical Research
Journal of Pharmacology, Pharmaceutics & Pharmacovigilance
Journal of Physical Medicine, Rehabilitation & Disabilities
Journal of Plant Science: Current Research
Journal of Psychiatry, Depression & Anxiety
Journal of Pulmonary Medicine & Respiratory Research
Journal of Practical & Professional Nursing
Journal of Reproductive Medicine, Gynaecology & Obstetrics
Journal of Stem Cells Research, Development & Therapy
Journal of Surgery: Current Trends & Innovations
Journal of Toxicology: Current Research
Journal of Translational Science and Research
Trends in Anatomy & Physiology
Journal of Vaccines Research & Vaccination
Journal of Virology & Antivirals
Archives of Surgery and Surgical Education
Sports Medicine and Injury Care Journal
International Journal of Case Reports and Therapeutic Studies
Journal of Ecology Research and Conservation Biology

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>