

Review Article

Relationship between Antioxidants and the Development of the Periodontal Disease

Rudrakshi C*, Prabhuji MLV, Parween S and Jyothisna S

Department of Periodontology, Krishnadevaraya College of Dental Sciences & Hospital, Hunasamaranahalli, Bangalore

Abstract

The role of oxidative stress linking the free radical damage at the cellular level causes premature aging, including periodontal disease. To overcome this, it is important to explore the role of host specifically with regard to antioxidant status along with conventional periodontal therapy. Periodontal counselling and supplementation may very well reduce inflammation and thereby enhance outcomes of conventional periodontal therapy. The purpose of this review is to summarize available research in the role of antioxidants as an adjunct to periodontal therapy.

Keywords: Antioxidants; Free radical; Inflammation; Periodontal therapy

Key Messages: Antioxidants have shown to have beneficial outcome when used as adjunct to traditional periodontal therapy. Implication of antioxidants has promising results in future.

Introduction

Periodontitis is a chronic inflammatory disease caused primarily by bacteria in dental plaque, affecting the supporting structures of the teeth. Specific periodontal pathogens such as the gram-negative anaerobic bacteria inhabiting within the sub-gingival plaque are associated with the progressive form of the disease. Although bacteria are the major etiological agents, the host immune response to these bacteria is of fundamental importance. Hence, it is evident that periodontitis is a multifactorial disease, affiliates with specific microorganisms, social and behavioral factors, genetic or epigenetic trait, all of which are modulated and controlled by the underlying immune and inflammatory responses of the host [1].

*Corresponding author: Rudrakshi C, Department of Periodontology, Krishnadevaraya College of Dental Sciences & Hospital, Hunasamaranahalli, Bangalore, Tel: +91 9448057407; E-mail: drudrakshi@rediffmail.com

Citation: Rudrakshi C, Prabhuji MLV, Parween S, Jyothisna S (2017) Relationship between Antioxidants and the Development of the Periodontal Disease. J Cytol Tissue Biol 4: 016.

Received: June 02, 2017; **Accepted:** August 12, 2017; **Published:** August 25, 2017

The recent focus on the progression of periodontitis is on the molecular aspects of host modulation. The discovery of the role of free radicals in chronic disease is as important as the discovery of the role of microorganism in inflammatory disease [2].

Free radicals & Reactive Oxygen Species (ROS) production is an essential component of the host response for immune system, prostaglandin biosynthesis, anti-bacterial biosynthesis and variety of insults like trauma/burns [3]. Besides, the physiological system, ROS generation are induced by several exogenous factors such as pollution, smoke, radiation, pesticides and drug consumption. Major producers of ROS are mitochondrial cytochrome P-450 reactions, peroxisomal fatty acid metabolism and NADPH oxidase activity [4]. An imbalance between the ROS production and anti-oxidant mechanism leads to oxidative stress. Oxidative stress has been associated with both onset of periodontal destruction and systemic inflammation [5,6].

Primary immune response against periodontal pathogens is elevated numbers of neutrophils seen in connective tissue, junctional epithelium (50%) and gingival crevicular fluid (90%) and it can cause loss of epithelial cell-cell attachment in junctional epithelium (>60%) leading to apical shift of junctional epithelium [7].

Neutrophil have several mechanisms for controlling bacterial invasion which includes both intracellular and extracellular oxidative and non-oxidative killing mechanisms [8]. When neutrophils and macrophages get stimulated by a phagocytic stimulus, produces a 'respiratory burst', which is characterized by an increase in oxygen consumption, activation of the Hexose-Monophosphate (HMP) shunt and generation of Free Radicals (FR), Reactive Oxygen Species (ROS) and their metabolic products. At sites of chronic inflammation (periodontium in case of periodontitis), there is considerable over production of FR and reactive species.

Research with scientific evidence confirmed that inflammation in the oral tissues especially that associated with periodontitis can be a factor in chronic illness such as cardio and cerebrovascular diseases [9,10], diabetes mellitus, rheumatoid arthritis, advance pregnancy outcomes and a growing list of other conditions such as cancer, psoriasis, chronic kidney disease and anemia. Several hypotheses proposed to describe the interlinking between periodontal diseases and systemic diseases. One of the explanations is imbalance between systemic oxidants and antioxidants [11-13]. "An antioxidant is any substance that, when present at low concentrations compared to those of an oxidizable substrate, significantly delays or prevents oxidation of that substrate" [14].

Hence the concept of inhibition of building up oxidative stress within cells through anti-oxidative therapy is implicated in inflammatory disorders and periodontitis along with gold standard therapy for periodontitis is through removal of subgingival biofilm through scaling and root planing has shown to be advantageous. The present article reviews on the comprehensive appraisal of the newer aspects of anti-oxidative therapy and it's applications in Periodontology (Figure 1).




Figure 1: Role of polymorphopneutrophils.

Reactive Oxygen Species (ROS)

Every day, numerous free radicals and ROS are produced in each cell by various exogenous and endogenous sources and these ROS are counteracted or balanced by anti-oxidants present within each cell, there by prevents oxidative stress build up. Reactive oxygen species and their sources are enumerated in table 1.

Recent literature postulate that, non-radicals which have the capability of radical transformation in the extra- and intracellular environment (singlet oxygen, hydrogen peroxide and hypochlorous acid) are grouped under reactive oxygen species category [15,16]. Tissue damage by ROS [5,15,17] (Table 2).

Free radicals Sources		Sources	
Oxygen derived free radical	Non- Oxygen derived free radical	Exogenous	Endogenous (byproducts)
			Metabolism
			Function
Superoxide (O ₂ ⁻)	Singlet oxygen (¹ O ₂)	Heat, Trauma	Cell metabolism and oxidative phosphorylation (electrons transport chain) By Osteoclasts at ruffled border during bone resorption. Oxidative pathway of phagocytosis is by neutrophils and other phagocytes.
Hydroxyl (OH [•])	Ozone (O ₃)	Ultrasound, Ultraviolet Light, Ozone, Smoking	
Hydroperoxyl (HOO [•])	Hypochlorous acid (HOCl)	Exhaust Fumes, Radiation, Infection, Excessive Exercise, Therapeutic Drugs	
Alkoxy (RO [•])	Hydrogen peroxide (H ₂ O ₂)		
Aryloxy (ArO [•])			
Arylperoxy (ArOO [•])			
Peroxy (ROO [•])			
Acyloxy (RCOO [•])			
Acylperoxy (RCOOO [•])			

Table 1: Reactive oxygen species and their sources.

ROS (Mainly Hydroxyl and peroxynitrate anion)	Target molecule	Example	Effects
	Protein	Gingival hyaluronic acid and Proteoglycans	<ul style="list-style-type: none"> Folding/unfolding, Fragmentation Protease degradation of the modified protein Formation of protein radicals and protein bond ROS polymerization of proteins Formation of stable end products (acetaldehydes)
	Lipid	Cell membrane (activation of cyclooxygenases and lipo-oxygenases pathways)	<ul style="list-style-type: none"> Peroxidation and formation of products these are bioactive molecules, Conjugated dienes Lipid Proxides, Aldehydes Volatile-hydrocarbons, Prostaglandin-E2(PG-E2) production eg: Malondialdehyde, Acrolein, Isoprostanes and Neuroprostanes
	DNA	Activating nuclear factor kB (NFkB)	<ul style="list-style-type: none"> Strand breaks, Base pair mutations, Deletions, Insertions, conversion of guanine to 8- hydroxyguanine, Nicking and Sequence amplifications. Stimulation of pro-inflammatory cytokine release by monocytes and macrophages
	Enzymes	Antiproteases such as α-1antitrypsin	Oxidation
Superoxide and hydrogen peroxide	Cells	Osteoclast activation	Bone resorption

Table 2: Tissue damage by ROS.


Halliwell established 4 criteria for causal relationship between ROS and disease [18];

1. ROS or the oxidative damage caused must be present at the site of injury.
2. The time course of ROS formation or the oxidative damage caused should occur before or at the same time as tissue injury.
3. Direct application of ROS over a relevant time course to tissues at concentrations found *in vivo* should reproduce damage similar to that observed in the diseased tissue.
4. Removing or inhibiting ROS formation should decrease tissue damage to an extent related to their antioxidant action *in vivo*.

Growing evidence exist in causal relationship between oxidative stress and periodontitis [19] (Table 3).

Targeted Periodontal tissue	Reaction of ROS	Effect
Ground substance	Depolymerization and Degradation (non-sulfated glycosaminoglycans are more susceptible than sulfated)	All these events leads to periodontal tissue destruction (Figure 2) and alveolar bone resorption
Collagen	Collagenolysis	
Monocytes and macrophages	Stimulation of excessive pro-inflammatory cytokine	
Lipid peroxidation	PG-E2 production leads to bone resorption	

Table 3: Reactive oxygen species on periodontal tissues.



Evidence suggesting causal relationship between ROS and periodontitis

It is more than a decade the physiological and pathological roles of ROS in the development of periodontitis have been studied

Chapple and Matthews postulated that periodontal tissue destruction results in increased production of ROS and increased ratio between elastase and lactoferrin by peripheral neutrophils [5].

Agnihotri have been reported that increased levels of ROS in smokers was responsible for excessive periodontal tissue destruction [21].

Patil et al., have reported that the severity of tissue destruction due to excessive ROS is more when periodontal disease is associated with type 2 DM, indicating that oxidative stress is common factor involved in tissue destruction [22].

Di Meo et al., suggested that low levels of ROS are beneficial, but excessive generation or antioxidant deficiency results in periodontal tissue destruction [23].

Liu et al., He et al., have shown the potential link between ROS and autophagy in periodontitis [16,24].

Antioxidants (AO)

The concept of inhibition of building up oxidative stress within cells through anti-oxidative therapy is implicated in inflammatory disorders and periodontitis. Antioxidants are agents which scavenge free radicals or otherwise reactive oxygen species, thereby preventing damage caused by them. Antioxidant therapy as adjunct to conventional treatments such as scaling and root planing has shown to be advantageous. Antioxidant can be classified in the following way (Table 4).

Sl.no	Based on	Examples
1	Mode of action	Preventive Suppress the formation of free radicals: Catalase, Glutathione peroxidase, and Serum transferrase Metal ion sequestrators- Albumin, Lactoferrin, Transferrin, Ceruloplasmin, Uric acid, Polyphenolic flavenoids Quenching of active oxygen: Superoxide dismutase, Carotenoids
		Scavenging (Chain breaking) Ascorbate, Carotenoids, Uric acid, Vitamin E, Bilirubin, Reduced glutathione & other
		2
	Intracellular	Superoxide dismutase enzymes-1& 2, Catalase, Glutathione peroxidase, DNA repair enzymes & Reduced glutathione.
	Extracellular	Superoxide dismutase enzyme-3, Slenium glutathione peroxidase, Reduced Glutathuione,
	Membrane associated	α-tocopherol
3	Solubility	Water soluble Haptoglobin, Ceruloplasmin, Albumin, Ascorbic Acid, Uric acid, Albumin, Polyphenolic flavenoids, Reduced glutathione & other thiols, Cysteine & Transferrins
		Lipid soluble α-tocopherol, Carotenoids, bilirubin, Ubiquinol and Vitamin A
4	Repair De novo enzymes	DNA repair enzymes, Protease, Lipase and Transferase
5	Structures they protect	DNA protective antioxidants Superoxide dismutase enzymes 1 and 2, Glutathione peroxidase, DNA repair enzymes [e.g., Poly(ADP-ribose) polymerase], Reduced glutathione, Cysteine
		Protein-protective antioxidants Sequestration of transition metals by preventative antioxidants Antioxidant enzymes
		Lipid-protective antioxidants α-Tocopherol (vitamin E), Ascorbate (vitamin C), Carotenoids (including retinol – vitamin A), Reduced ubiquinone, Reduced glutathione, Glutathione peroxidase, Bilirubin
6	By their origin	Exogenous antioxidants Carotenoids, Ascorbic acid, Tocopherols (a, b, c, d), Polyphenols (e.g. Flavenoids, Catechins such as Epigallocatechin-gallate), Folic acid, Cysteine
		Endogenous antioxidants Catalase, Superoxide dismutase, Glutathione peroxidase, Glutathione-S-transferase, Reduced glutathione, Ceruloplasmin, Transferrin, Ferritin, Glycosylases, Peroxisomes, Proteases
		Synthetic N-acetylcysteine, Penicillinamine, Tetracyclines

Table 4: Anti-oxidant classification.

Antioxidant enriched diet and periodontal diseases [25]

Recent evidence has demonstrated a significant role for oxidative stress in promoting bone resorption via activation of certain transcription factors (Fox Os, which decreases wnt signalling), Modulated by insulin resistance and increasing age [26]. Antioxidant micronutrients combat such pro-inflammatory cascades through modulation of oxidative stress by directly scavenging Reactive Oxygen Species (ROS) and also by down-regulation of some redox-sensitive pro-inflammatory gene transcription factors such as nuclea factor-kappa B and activator protein-1 while up regulating anti-inflammatory gene transcription factors such as nuclear factor [5]. Following are some of the examples of antioxidant diets.

Green Tea

Green tea is a non-fermented product of tea (*Camellina sinensis*) leaves that is consumed as a beverage worldwide. Green tea extract is documented to have antibacterial, antioxidant, anti-inflammatory and anticarcinogenic properties. It is a rich source of flavonoids, mainly catechins. The four major catechins include

- Epigallocatechin-3-gallate (59%)
- Epigallocatechin (19%)
- Epicatechin-3-gallate (13.6%)
- Epicatechin

Local drug delivery of green tea extract has also shown promising results in treating periodontal diseases [27]. Dentifrice and mouthwashes can be contemplated as vehicles for self-application of chemotherapeutics. Recently, the addition of green tea catechins to dentifrice has shown reduction of periodontal inflammation in the rat model, by decreasing gingival oxidative stress and expression of pro-inflammatory cytokines [28].

Hrishi et al., conducted a study to evaluate the effect of adjunctive use of green tea dentifrice in periodontitis patients and concluded that green tea dentifrice may serve as a beneficial adjunct to non-surgical periodontal therapy as it showed greater reduction of gingival inflammation and improved periodontal parameters on comparison with fluoride-triclosan dentifrice [29].

Grape Seed Extracts (GSE)

Grape seed extract is a naturally occurring polyphenolic compound obtained from seeds of *Vitis vinifera*. It possesses a wide range of biological activities such as immunomodulator agent, antioxidant, anticarcinogenic, and anti-inflammatory effects [30,31]. The immunomodulator effect is particularly due to its proanthocyanidin content. GSE may be beneficial for the treatment of inflammation associated with bone destruction as it may strongly inhibit osteoclast differentiation, reduce osteoclast activity, and stimulate bone formation through its positive action on osteoblast differentiation [32]. The anti-inflammatory effect is modulated by calibrating the delicate balance between pro-inflammatory and anti-inflammatory cytokines through regulating their release and gene expression delicate balance between pro-inflammatory and anti-inflammatory cytokines through regulating their release and gene expression [33].

Ozden FO et al., investigated the effects of GSE application on periodontium before and after ligature induced experimental periodontitis [34], using histological and immunohistochemical analyses. Histopathological findings showed improvements in the inhibition of periodontal inflammation and destruction following GSE intake.

The bark contains magnolol and honokiol, two polyphenolic compounds that have been demonstrated as peroxisome proliferator-activated receptor gamma (PPAR gamma) agonists and gamma-aminobutyric acid modulators. Magnolol extracted from *Magnolia officinalis* is widely used in oriental medicine. Preclinical studies have evaluated their various potential applications including antioxidant, anti-inflammatory, antitumor and antimicrobial properties [35].

Magnolol has been reported to have a potent anti-inflammatory activity via inhibition of proinflammatory cytokine, ROS formation, iNOS, COX-2 expression, and nuclear factor-kappaB (NF-kappaB) activation, a key transcription factor regulating inflammation, in LPS-induced inflammatory disease [36,37]. Furthermore, magnolol exerts a marked antimicrobial activity against periodontopathic bacteria and activates osteoblast function [38]. Magnolol may be used as a potential candidate for treating periodontitis.

Lu SH et al., concluded in their study that magnolol significantly ameliorates the alveolar bone loss in ligature-induced experimental periodontitis by suppressing periodontopathic microorganism accumulation [39], NF-κB-mediated inflammatory mediator synthesis, RANKL formation, and osteoclastogenesis. These activities support that magnolol is a potential agent to treat periodontal disease.

Melatonin

Melatonin (N-acetyl-5-methoxy tryptamine) is a substance secreted by multiple organs including the pineal gland, retina, bone marrow, the gastro-intestinal track and the immune system. Because it is lipophilic, it can reach all body cells and cellular components rapidly [40]. In addition to its scavenger effects, melatonin indirectly stimulates various antioxidant enzymes [41,42]. A quality that makes melatonin superior to other antioxidants is that it does not become pro-oxidative by losing an electron during interactions with free radicals; i.e., it is not a potential pro-oxidant [43].

Local and systemic administration of melatonin in rats with lipopolysaccharide-induced periodontitis reduced the level of enzymes (such as serum aspartate aminotransferase, alanine transaminase and blood urea nitrogen) significantly compared with rats in the control group [42,44]. Similarly, locally administered melatonin significantly reduced bone resorption compared with rats receiving no treatment. These studies suggested that topical administration of melatonin can be used as an adjunct to conventional treatment protocols such as scaling, root planing, and surgical debridement to improve the outcomes of periodontal therapy.

Propolis Extract

Propolis, sometimes called bee glue, is a natural resinous substance collected by honey bees (*Apis mellifera L.*) from plant buds and bark exudates. Bio-flavonols are the key contributors to propolis' properties. Propolis is found to have strong inhibitory effects on at least 21 species of bacteria, 9 species of fungi, 3 species of protozoa, and a wide range of viruses.

A number of studies have presented evidence that propolis has strong hepatoprotective, antitumor, antioxidative, antimicrobial and anti-inflammatory properties. Coutinho A in the year 2012 in their clinical study demonstrated the benefits provided by propolis extract and indicated that it should be considered for use as an adjuvant to scaling and root planing [45-48].

Salivary Antioxidant Status

Saliva is rich in antioxidants, mainly uric acid accounting for more than 70% of the total antioxidant activity of resting and stimulated saliva from both healthy and periodontally compromised subjects. Other antioxidants present in lesser concentration than uric acid are albumin, ascorbate and glutathione [49]. Albumin concentration is comparatively low, about 10mM, apparently in the same range of that of ascorbic acid [49,50]. Accounting for the 5-10% of antioxidant activity in saliva are the traces of other antioxidants such as transferrin, lactoferrin and caeruloplasmin. They are capable of binding metal ions are found in both saliva and GCF [51].

Antioxidant status in gingival crevicular fluid

An excess of reactive oxygen species and depletion of antioxidant levels in gingival crevicular fluid are responsible for the chronic local activation of periodontal inflammation and tissue destruction [5,52]. Gingival crevicular fluid probably contains a locally derived antioxidant from local three sources such as plaque bacteria, neutrophils and crevicular epithelium [53].

Significantly more intense levels of superoxide dismutases in gingival crevicular fluid comprise the most important antioxidant enzyme defense system against reactive oxygen species.

The reduced and oxidised glutathione concentrations in the gingival crevicular fluid are significantly lower in patients with chronic periodontitis, supporting that the glutathione levels in gingival crevicular fluid can play an important role in the pathogenesis of periodontitis [54].

Total antioxidant capacity

It is the measure of the antioxidant capacity of all antioxidants in a biological sample rather than the antioxidant capacity of a single compound. Total Antioxidant Capacity (TAC) of saliva has been measured by only 3 methods using 3 different biochemical techniques [55];


- Spectrophotometric assay
- Enhanced chemiluminescence assay
- Cyclic voltammetry assay

Salivary TAC decreases or does not changes in periodontal disease.

Mechanism of Action [5,15]



Superoxide Dismutase (SOD)

It is an anti-oxidative enzyme, which has been localized within human periodontal ligament and represent an vital defense mechanism for gingival fibroblasts against superoxide ions by catalyzing and accelerating the reaction (approximately 10,000 times) from superoxide ion to oxygen and less reactive H_2O_2 (Figure 3).




In humans SOD exists in 3 forms (Figures 4 and 5)

- Cytosolic Cu/Zn-SOD,
- Mitochondrial Mn-SOD (major role)
- Extracellular SOD (EC-SOD)




Catalase (CAT)

It is heme bound iron anti-oxidative enzyme, majorly located in peroxisomes. It protects the cells from hydrogen peroxide by converting it into water (Figure 6).



Reduced Glutathione (GSH)

GSH is an essential antioxidant enzyme, which AIDs as A- Anti-oxidant, I-Immunobooster, D- Detoxifier. It regulates IL-2 dependent T lymphocyte proliferation (Figure 7).



Glutathione Peroxidase

It is a selenium containing peroxidase which reduces variety of hydroperoxides, H₂O₂ and lipids, there by protects the cell particularly against low levels of oxidative stress. It has 5 types of isoenzymes, these are GPX1, GPX2, GPX3, GPX4, and GPX5.

Uric acid

Moore et al., in 1994 proposed that uric acid is responsible for 70% antioxidants capacity of saliva and it is one of the major antioxidant present in saliva. Diab-Ladki in 2003 [56] found decreased concentrations of uric acid, ascorbic acid and albumin in saliva of periodontitis patients.

Ascorbic acid (vitamin C)

It is a water soluble vitamin, function as antioxidant by scavenging water-soluble peroxy radicals, scavenging superoxide and perhydroxyl radicals, prevention of damage mediated by hydroxyl radicals on uric acid, scavenger of hypochlorous acid, decreases heme breakdown and subsequent Fe²⁺ release thereby preventing Fenton reactions, scavenger of singlet oxygen and hydroxyl radicals, re-forms a-tocopherol from its radical and also protects against ROS-release from cigarette smoke.

Conclusion

Following conclusion can be drawn with respect to the role of reactive oxygen species and antioxidants in periodontitis

1. Oxidative stress and reactive oxygen species appear to play a significant role in the patho-physiology of periodontal diseases.
2. Adjunctive use of antioxidants with traditional therapies should be considered to improve the periodontal treatment outcome.
3. This review presents with evidence within the biomedical literature, provides exciting new opportunities for future development of host modulation therapies in periodontology.

Future Research Directions

The results of the review may have relevance for the development of treatment protocols. The impact of the combination of periodontal therapy with antioxidants in terms of antioxidant/oxidative stress parameters requires further investigation with longer follow.

References

1. Lamster IB, Novak MJ (1992) Host mediators in gingival crevicular fluid: Implications for the pathogenesis of periodontal disease. *Crit Rev Oral Bio Med* 3: 31-60.
2. Bray TM (1999) Antioxidants and Oxidative Stress in Health and Disease: Introduction. *Proc Soc Exp Biol Med* 222: 195.
3. Pham-Huy LA, He H, Pham-Huy C (2008) Free radicals, antioxidants in disease and health. *Int J Biomed Sci* 4: 89-96.
4. Downey GP, Fukushima T, Fialkow L (1995) Signaling mechanisms in human neutrophils. *Curr Opin Hematol* 2: 76-88.
5. Chapple IL, Matthews JB (2007) The role of reactive oxygen and antioxidant species in periodontal tissue destruction. *Periodontol* 2000 43: 160-232.
6. Basu S, Helmersson J, Jarosinska D, Sällsten G, Mazzolai B, et al. (2009) Regulatory factors of basal F(2)-isoprostane formation: population, age, gender and smoking habits in humans. *Free Radic Res* 43: 85-91.
7. Cekici A, Kantarci A, Hasturk H, Van Dyke TE (2014) Inflammatory and immune pathways in the pathogenesis of periodontal disease. *Periodontol* 2000 64: 57-80.
8. Dennison DK, Van Dyke TE (1997) The acute inflammatory response and the role of phagocytic cells in periodontal health and disease. *Periodontol* 2000 14: 54-78.
9. Eke PI, Dye BA, Wei L, Slade GD, Thornton-Evans GO, et al. (2015) Update on prevalence of periodontitis in adults in the United States: NHANES 2009 to 2012. *J Periodontol* 86: 611-622.
10. Engebretson S, Kocher T (2013) Evidence that periodontal treatment improves diabetes outcomes: A systematic review and meta-analysis. *J Clin Periodontol* 40: 153-163.
11. Jin LJ, Lamster IB, Greenspan JS, Pitts NB, Scully C, et al. (2016) Global burden of oral diseases: Emerging concepts, management and interplay with systemic health. *Oral Dis* 22: 609-619.
12. Buset SL, Walter C, Friedmann A, Weiger R, Borgnakke WS, et al. (2016) Are periodontal diseases really silent? A systematic review of their effect on quality of life. *J Clin Periodontol* 43: 333-344.
13. Jeffcoat MK, Jeffcoat RL, Gladowski PA, Bramson JB, Blum JJ (2014) Impact of periodontal therapy on general health: Evidence from insurance data for five systemic conditions. *Am J Prev Med* 47: 166-174.
14. Halliwell B (1997) Antioxidants: The basics--what they are and how to evaluate them. *Adv Pharmacol* 38: 3-20.
15. Sree SL, Mythili R (2011) Antioxidants in Periodontal Diseases : A Review. *Indian J Multidiscip Dent* 1: 140-146.
16. Chengcheng Liu, Longyi Mo, Yulong Niu, Xin Li, Xuedong Zhou, et al. (2017) The Role of Reactive Oxygen Species and Autophagy in Periodontitis and Their Potential Linkage. *Front Physiol* 8: 439.
17. Dahiya P, Kamal R, Gupta R, Bhardwaj R, Chaudhary K, et al. (2013) Reactive oxygen species in periodontitis. *J Indian Soc Periodontol* 17: 411-416.
18. Halliwell B (1989) Free radicals, reactive oxygen species and human disease: A critical evaluation with special reference to atherosclerosis. *Br J Exp Pathol* 70: 737-757.
19. Pendyala G, Thomas B, Kumari S (2008) The challenge of antioxidants to free radicals in periodontitis. *J Indian Soc Periodontol* 12: 79-83.
20. Sharma A, Wati S, Sharma S (2011) Reactive oxygen species and antioxidants in periodontics : A Review. *Int J Dent Clin* 3: 44-47.
21. Agnihotri R, Pandurang P, Kamath SU, Goyal R, Ballal S, et al. (2009) Association of cigarette smoking with superoxide dismutase enzyme levels in subjects with chronic periodontitis. *J Periodontol* 80: 657-662.
22. Patil VS, Patil VP, Gokhale N, Acharya A, Kangokar P (2016) Chronic Periodontitis in Type 2 Diabetes Mellitus: Oxidative Stress as a Common Factor in Periodontal Tissue Injury. *J Clin Diagn Res* 10: 12-16.
23. Di Meo S, Reed TT, Venditti P, Victor VM (2016) Role of ROS and RNS Sources in Physiological and Pathological Conditions. *Oxid Med Cell Longev* 2016: 1245049.
24. He ZJ, Zhu FY, Li SS, Zhong L, Tan HY, et al. (2017) Inhibiting ROS-NF-κB-dependent autophagy enhanced brazilin-induced apoptosis in head and neck squamous cell carcinoma. *Food Chem Toxicol* 101: 55-66.
25. Muniz FW, Nogueira SB, Mendes FL, Rösing CK, Moreira MM, et al. (2015) The impact of antioxidant agents complimentary to periodontal therapy on oxidative stress and periodontal outcomes: A systematic review. *Arch Oral Biol* 60: 1203-1214.
26. Galli C, Passeri G, Macaluso GM (2011) FoxOs, Wnts and oxidative stress-induced bone loss: new players in the periodontitis arena? *J Periodontol Res* 46: 397-406.

27. Hirasawa M, Takada K, Makimura M, Otake S (2002) Improvement of periodontal status by green tea catechin using a local delivery system: a clinical pilot study. *J Periodontol Res* 37: 433-438.
28. Kudva P, Tabasum ST, Shekhawat NK (2011) Effect of green tea catechin, a local drug delivery system as an adjunct to scaling and root planing in chronic periodontitis patients: A clinicomicrobiological study. *J Indian Soc Periodontol* 15: 39-45.
29. Hrishi TS, Kundapur PP, Naha A, Thomas BS, Kamath S, et al. (2016) Effect of adjunctive use of green tea dentifrice in periodontitis patients - A randomized controlled pilot study. *Int J Dent Hyg* 14: 178-183.
30. Baydar NG, Sagdic O, Ozkan G, Cetin S (2006) Determination of antibacterial effects and total phenolic contents of grape (*Vitis vinifera* L.) seed extracts. *Int J Food Sci Tech* 41: 799-804.
31. Zhang XY, Bai DC, Wu YJ, Li WG, Liu NF (2005) Proanthocyanidin from grape seeds enhances anti-tumor effect of doxorubicin both in vitro and in vivo. *Pharmazie* 60: 533-538.
32. Park JS, Park MK, Oh HJ, Woo YJ, Lim MA, et al. (2012) Grape-seed proanthocyanidin extract as suppressors of bone destruction in inflammatory autoimmune arthritis. *PLoS One* 7: 51377.
33. Ahmad SF, Zoheir KM, Abdel-Hamied HE, Attia SM, Bakheet SA, et al. (2014) Grape seed proanthocyanidin extract protects against carrageenan-induced lung inflammation in mice through reduction of pro-inflammatory markers and chemokine expressions. *Inflammation* 37: 500-511.
34. Özden FO, Sakallioğlu EE, Sakallioğlu U, Ayas B, Erişgin Z (2017) Effects of grape seed extract on periodontal disease: an experimental study in rats. *J Appl Oral Sci* 25: 121-129.
35. Shen JL, Man KM, Huang PH, Chen WC, Chen DC, et al. (2010) Honokiol and magnolol as multifunctional antioxidative molecules for dermatologic disorders. *Molecules* 15: 6452-6465.
36. Yang TC, Zhang SW, Sun LN, Wang H, Ren AM (2008) Magnolol attenuates sepsis-induced gastrointestinal dysmotility in rats by modulating inflammatory mediators. *World J Gastroenterol* 14: 7353-7360.
37. Tsai YC, Cheng PY, Kung CW, Peng YJ, Ke TH, et al. (2010) Beneficial effects of magnolol in a rodent model of endotoxin shock. *Eur J Pharmacol* 641: 67-73.
38. Kwak EJ, Lee YS, Choi EM (2012) Effect of Magnolol on the Function of Osteoblastic MC3T3-E1 Cells. *Mediators of Inflammation* 2012: 829650.
39. Lu SH, Huang RY, Chou TC (2013) Magnolol ameliorates ligature-induced periodontitis in rats and osteoclastogenesis: In vivo and in vitro study. *Evid Based Complement Alternat Med* 2013: 634095.
40. Menendez-Pelaez A, Poeggeler B, Reiter RJ, Barlow-Walden L, Pablos MI, et al. (1993) Nuclear localization of melatonin in different mammalian tissues: Immunocytochemical and radioimmunoassay evidence. *J Cell Biochem* 53: 373-382.
41. Kara A, Akman S, Ozkanlar S, Tozoglu U, Kalkan Y, et al. (2013) Immune modulatory and antioxidant effects of melatonin in experimental periodontitis in rats. *Free Radic Biol Med* 55: 21-26.
42. Arabacı T, Kermen E, Özkanlar S, Köse O, Kara A, et al. (2015) Therapeutic effects of melatonin on alveolar bone resorption after experimental periodontitis in rats: A biochemical and immunohistochemical study. *J Periodontol* 86: 874-881.
43. Czesnikiewicz-Guzik M, Konturek SJ, Loster B, Wisniewska G, Majewski S (2007) Melatonin and its role in oxidative stress related diseases of oral cavity. *J Physiol Pharmacol* 58: 5-19.
44. Gulle K, Akpolat M, Kurcer Z, Cengiz MI, Baba F, et al. (2014) Multi-organ injuries caused by lipopolysaccharide-induced periodontal inflammation in rats: role of melatonin. *J Periodontol Res* 49: 736-741.
45. Banskota AH, Tezuka Y, Kadota S (2001) Recent progress in pharmacological research of propolis. *Phytother Res* 15: 561-571.
46. Sforcin JM (2007) Propolis and immune system: A review. *J Ethnopharmacol* 113: 1-14.
47. Martos VM, Navajas RY, Fernandez LJ, Alvarez JA (2008) Functional properties of honey, propolis, and royal jelly. *J Food Sci* 73: 117-124.
48. Coutinho A (2012) Honeybee propolis extract in periodontal treatment: a clinical and microbiological study of propolis in periodontal treatment. *Indian J Dent Res* 23: 294.
49. Phaniendra A, Jestadi DB, Periyasamy L (2015) Free radicals: Properties, sources, targets, and their implication in various diseases. *Indian J Clin Biochem* 30: 11-26.
50. Moore S, Calder KA, Miller NJ, Rice-Evans CA (1994) Antioxidant activity of saliva and periodontal disease. *Free Radic Res* 21: 417-425.
51. Lynch E, Sheerin A, Claxton AW, Atherton MD, Rhodes CJ, et al. (1997) Multicomponent spectroscopic investigations of salivary antioxidant consumption by an oral rinse preparation containing the stable free radical species chlorine dioxide (ClO₂). *Free Radic Res* 26: 209-234.
52. Chapple IL (1997) Reactive oxygen species and antioxidants in inflammatory diseases. *J Clin Periodontol* 24: 287-296.
53. Chapple ILC, Brock G, Eftimiadi C, Matthews JB (2002) Glutathione in gingival crevicular fluid and its relation to local antioxidant capacity in periodontal health and disease. *Mol Pathol* 55: 367-373.
54. Öngöz DF, Bozkurt DS, Ballı U, Avci B, Durmuşlar MC, et al. (2016) Glutathione levels in plasma, saliva and gingival crevicular fluid after periodontal therapy in obese and normal weight individuals. *J Periodont Res* 51: 726-734.
55. Kusano C, Ferrari B (2008) Total Antioxidant Capacity: A biomarker in biomedical and nutritional studies. *J Mol Cell Biol* 7: 1-15.
56. Diab-Ladki R, Pellat B, Chahine R (2003) Decrease in the total antioxidant activity of saliva in patients with periodontal diseases. *Clin Oral Investig* 7: 103-107.

