

## Research Article

# Influence of Organic Matrix and Cations on Bio-Methane Yield with Anthracite Methanogenic Consortium

Dong Xiao<sup>1</sup>, Yong Hou<sup>2\*</sup>, Enyuan Wang<sup>3</sup> and Yidong Zhang<sup>1</sup><sup>1</sup>State Key Laboratory of Coal Resources and Safe Mining, China University of Mining and Technology, Xuzhou, Jiangsu, China<sup>2</sup>Xuzhou Hongqiao New Energy Group Co., Ltd., Xuzhou, Jiangsu, China<sup>3</sup>School of Safety Engineering, China University of Mining and Technology, Xuzhou, Jiangsu, China

## Abstract

Organic compounds fermentation of coal has been used to generate secondary biogenic gas and enhance gas reservoirs in coal bed. To enhance the bio-degradation process, culture nutrition plays an important role in remediating the nutritional deficiency of the coal seam. The influence of bio-methane yield with organic inputs and cation concentrations was examined. Research of organic matrix influence revealed that the traditional organic material except yeast extract should forbid, and the input of yeast extract should limit at 1.00g/L also. Further, the study demonstrated that the ion concentration of sodium, potassium, magnesium, calcium and ammonia nitrogen also influenced methane and carbon dioxide yields. And the optimize concentrations for  $\text{Ca}^{2+}$ ,  $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Mg}^{2+}$  were 5.1, 1.7, 23 and 1.3 mmol/L. The  $\text{Mg}^{2+}$  was particularly sensitive in inhibiting  $\text{CH}_4$  metabolism processes largely for gas-coal methanogenic consortium.

**Keywords:** Anaerobic bacteria; Bio-methane; Coal; Culture media; Organic matrix

## Introduction

The demand for gas has seen a tremendous increase in the last 10 years [1]. In particular, coal-bed methane enhance with biotechnology represents an ideal method to increase methane productivity from oil deposits and coal seams using underground anaerobic digestion

\*Corresponding author: Yong Hou, Xuzhou Hongqiao New Energy Group Co., Ltd., Xuzhou, Jiangsu, China, China, Tel: 86 18626026321; E-mail: hy\_9999@126.com

**Citation:** Xiao D, Hou Y, Wang E, Zhang Y (2020) Influence of Organic Matrix and Cations on Bio-Methane Yield with Anthracite Methanogenic Consortium. Adv Microb Res 4: 011.

**Received:** April 28, 2020; **Accepted:** May 13, 2020; **Published:** May 20, 2020

**Copyright:** © 2020 Xiao D, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

processing [2,3]. With the addition of adequate nutrient supplements, methanogenic consortium has the potential to degrade organic compounds in coal into methane. In this way, gas reserves and productivity could be enhanced along with secondary biogenic gas generation [4-6].

Previous studies involved with the methanogenic microbial traits and fermentation characteristics for bio-methane generation have focused on the processes of bio-degradation by microbial communities in coal seam [2,7,8]. Anion research identified that methane biosynthesis process could active where coal bed waters exhibit relatively low salinities ( $<2 \text{ mol/L Cl}$ ) and low  $\text{SO}_4^{2-}$  concentrations ( $<10 \text{ mmol/L}$ ) [9]. Nutrients supply by meteoric water plays an important role in methanogenic microbial metabolism [10,11]. However, nutritional deficiency exists in most coal seams. Most organic compounds in coal cannot be readily used as nutrients for methanogenic microbes. In addition to the inherent elements present in coal beds, supplemental nutrients are important to remediate the nutritional deficiency and promote microbial growth [12,13]. If concentrations of supplemental nutrients are excessive, the methanogenic consortium will be adversely influenced [14]. The goal of this report to identify the optimal nutrient dose for microbial community culture. Methane and carbon dioxide yield rate were the indicators used to assess the results of experiments.

## Materials and methods

### Materials

The coal and coal bed water samples used in this study were collected from Laohutai Fm 103<sup>#</sup> coal bed located in Shenyang province of China (GPS coordinates 41.830256, 123.957639). The depth of the Fm 103<sup>#</sup> gas-coal sample was 550 meters and the thickness of the coal seam was 6-18 meters. Laohutai Mine, in the east part of Fushun near Shenyang, Liaoning province, opened in 1901. The mine has 55.6 million tons of recoverable coal reserves until 2004 [15]. The samples were collected by the State Key Laboratory of Coal Resources and Safety Mining. The field studies did not involve endangered or protected species, and only small sample quantities were used. And no specific permissions were required for these locations sample collection and research.

The coal samples were obtained from a newly exposed coal seam in the heading face by the channel sampling method following ASTM D 4596-86 with nitrogen protection. Two sample points were selected from the Fm 103<sup>#</sup> coal seam. These samples were sealed in gas-tight steel canisters and set with gas-tight valves (manufactured by J&D Technologies) which enabled inert gas to flush the canister and protected with nitrogen immediately. The coal to be used for the microbe community source was crushed into small pieces (10-15 mm in diameter) at an aseptic bench (manufactured by Jing Xue Technologies). Nitrogen protection was used to maintain the coal in a continuous anaerobic environment. The sample was sealed in a sterilized gas desorption canister (manufactured by J&D Technologies) to desorb the absorbed gas until no gas desorbed in atmospheric pressure at 25°C [16].

The formation of water samples from the Fm 103<sup>#</sup> coal bed were collected in the same coal-bed in sterile glass bottles (manufactured by Fisher Scientific) which were filled to overflow to prevent oxygen ingress. Coal bed water ( $\leq 24$  hours after collection) was autoclaved at 121°C for 45 minutes and sealed in sterile glass bottles. Argon was infused to seal the top space of the bottles. The coalbed water samples were stored at 4°C for no more than 3 days.

## Culture

Five different nutrition culture media were tested in these experiments: MAC-1, MAC-2, MAC-3, MAC-4 and MAC-5 (MAC is the abbreviation of “the influence test of organic matrix and cautions”) [4,17]. The final concentrations of the compounds (g/L) were as list in table 1 and 0.1 mL/L of resazurin was added as an oxygen indicator. Yeast extract was produced by Fisher BioReagents, and other chemicals were supplied by Acros Organics.

Culture Media	Sodium Acetate	Glucose	Beef Extract	Yeast Extract	Common Concentrations
MAC-1	2.00	3.00	3.00	2.00	NH <sub>4</sub> Cl, 1.00; MgCl <sub>2</sub> ·7H <sub>2</sub> O, 1.00; KH <sub>2</sub> PO <sub>4</sub> , 0.40; KCl, 0.50; NaHCO <sub>3</sub> , 1.00; L-Cysteine Hydrochloride Monohydrate, 0.45
MAC-2	2.00	1.00	1.50	1.00	
MAC-3	0.00	1.00	1.50	1.00	
MAC-4	0.00	0.00	0.00	1.00	
MAC-5	0.00	0.00	0.00	0.00	

**Table 1:** The MAC1-MAC5 nutrition culture media concentrations (g/L).

The distilled water was autoclaved at 121°C for 45 minutes with dissolved oxygen removed. Nutrition medium was prepared in a 500 mL flask with distilled water. The medium was mixed using a magnetic stirrer for 2 hours at 60°C at an aseptic bench and then combined with an equal volume of coal bed water (volume ratio1:1) for another hour at room temperature. Nitrogen protection was used to maintain nutrition in a continuous anaerobic environment throughout the entire experiment. The final pH was maintained at 6.0 for all nutrition cultures. Control samples of 100 mL MAC-1, MAC-2, MAC-3, MAC-4 and MAC-5 were sealed in separate sterile glass bottles and stored at -40°C Argon was infused to seal the top space of bottles.

Anaerobic conditions were ensured in flasks using a gas-replacement method. This gas replacing process was monitored in real time with a carbon dioxide monitor system (manufactured by E2V). For each experiment 50.00±1 g of the coal sample and 500.00 mL of medium were used. Nitrogen was used to seal the upper space of the flask at the beginning of the experiment. The flasks were placed in an incubation shaker at 35°C and were agitated at 80 rpm to maximize the coal-liquid mass transfer rates. Twelve parallel tests were designed for each nutrition group which were cultured for 40 days under identical conditions.

The control experiments were performed without coal sample, which named MAC-1\*, MAC-2\*, MAC-3\*, MAC-4\* and MAC-5\*, to identify whether exogenous organic has the possibility to supply extra carbon to enhance the bio-methane yield. For each control experiment of each nutrition group 100 mL of cultured medium and 400 mL new medium was used. And cultured for 40 days in the same condition as above.

## The cation orthogonal analysis

The caution orthogonal analysis was based on consumption of cation elements in bio-degradation processes, according to the L<sub>16</sub>(4<sup>5</sup>) orthogonal table which includes Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup> and Mg<sup>2+</sup>. The lowest ion level was based on the MAC-4 medium while the highest ion level was based upon that of the East China Sea ion concentration. A quantity of 50.00±1 g of Fm 103<sup>#</sup> gas-coal sample was used in each experiment. The upper space of the flasks was sealed with nitrogen. The flasks were placed on an incubation shaker at 35°C and agitated at 80 rpm to maximize coal-liquid mass transfer rates. Caution orthogonal experiments were of 40 days of culture.

## Gas analysis

Gas samples were obtained with a 50-μL gas syringe. Methane and carbon dioxide analyses were performed using an Agilent 7890 A gas chromatograph (manufactured by Agilent). The nitrogen (carrier gas) flow rate was set at 1.00 mL/min. The injection port was maintained at 150°C with the oven temperature set at 25°C and the Thermal Conductivity Detector (TCD) at 200°C. Retention times for methane were 3.76 minutes and 5.0 minutes for carbon dioxide. Calibration standards consisted of 40% methane, 20% carbon dioxide, 10% hydrogen and 30% nitrogen, which were injected at atmospheric pressure to generate the calibration plot.

## Nutrient metabolism analysis

Nutrient ion concentrations were analyzed using an HC-800 ionization analyzer (manufactured by Histrong Technologies). The main indicators included fluoride, chloride, nitrate and nitrite nitrogen, phosphate, sulfate, carbonate, bicarbonate, ammonia nitrogen, sodium, potassium, magnesium, calcium, pH, water hardness and total alkalinity.

## Results


### The nutrient abundance influence for bio-degradation of coal

Five different organic compound nutrient concentrations designated as MAC-1, MAC-2, MAC-3, MAC-4 and MAC-5 were assessed in this study. In these medium, Sodium Acetate is a collective medium, as carbon resource, for methanogenic normally [18]. Glucose is the medium for hydrogen-producing acetogens. Beef Extract is a mixture of peptides and amino acids, nucleotide fractions, organic acids, minerals and some vitamins. It is often used to supply carbon and nitrogen sources. Yeast extract contains a mixture of amino acids, peptides, water soluble vitamins and carbohydrates. And it is often used in culture media [19]. The rank order of organic concentrations were MAC-1 > MAC-2 > MAC-3 > MAC-4 > MAC-5.

Coal bed methanogenic groups are comprised of a variety of microbial types. With regard to methanogens, only a limited number of simple carbon compounds such as CO<sub>2</sub> or acetate can serve as substrates. For the conversion of complex organic compounds to methane, fermentative and acetogenic bacteria are required. Thus they group an interactive methanogenic consortium [20]. In the process of gas-coal bio-degradation, the capacity for fermentative bacteria to hydrolyze and ferment the organic compounds of coal plays an important role, and CO<sub>2</sub> is the main compound of gas productions. Beef extract in medium supplied the extra carbon and nitrogen for bacteria. For the acetogenic bacteria fermentation process, the long chain fatty

acids and sugar degrade to form acetate, CO<sub>2</sub> and H<sub>2</sub> [18]. Glucose is the medium to enhance the acetogenic bacteria fermentation. And methanogens yield CH<sub>4</sub> with CO<sub>2</sub> and H<sub>2</sub> or acetate. Sodium Acetate was introduced to raise the acetate supply for methanogens [21].

If consortia in a good balance, most CO<sub>2</sub> and H<sub>2</sub> will be used to format CH<sub>4</sub>. And the concentration of CH<sub>4</sub> should be high; meanwhile, the concentration of CO<sub>2</sub> and H<sub>2</sub> should keep in low. So the yield gas concentration with microbial fermentation process could reflect the bacteria balance conditions. The presence of sufficient organic nutrients in the medium could promote fermentative and acetogenic bacteria flourish and improve methanogen nutrient generation. However, if excessive amounts of organic nutrients are contained within the medium, which like beef extract and glucose, the high propagation of fermentative or acetogenic bacteria would break the microbial balance, and carbon dioxide yield rate could be enhanced, meanwhile, methanogenesis could be inhibited. This phenomenon showed in MAC-1, MAC-2 and MAC-3 culture experiments (Figure 1).



The results confirmed that MAC-4 was the most effective medium for enhancing the bio-methane generation rate. 1 g/L yeast extract provided the best concentration of organics amino acids, peptides, and water-soluble vitamins to optimize the microbial fermentation. The maximal methane concentrations of the MAC-4 culture group achieved 23.62% (Figure 1). It was 4 times higher than that of the MAC-2, MAC-3, MAC-5 groups on average. The MAC-4\* control experiment and MAC-5 group verified that the anthracite is important to provide carbon for bio-methane yield (Figure 1).

Figure A is the methane concentration changes with culture days. And figure B is the control experiment without coal sample supply in experiment. Figure C is the carbon dioxide concentration changes with culture days. And figure D is the control experiment without coal sample supply in experiment. Methane is the main factor to identify the methanogens activity. And carbon dioxide is an important factor to identify the fermentative and acetogenic bacteria activity. The methane yield rate would high, only if the fermentative and

acetogenic bacteria activity in a limited condition. Organic material could enhance the fermentative and acetogenic bacteria. However, if the activity of fermentative and acetogenic bacteria is too high, it would inhibit the methanogens. The microbial group will in a good balance when achieved the highest methane yield and the best ratio of methane and carbon dioxide. Thus the MAC-4 medium culture group fit for the requirements.

The high carbon dioxide and low methane concentration data obtained from MAC-1 and MAC-2 media demonstrate that hydrolytic bacterial had enhanced with an abundance of organic substrate supplement, however, methane biosynthesis tended to inhibit in culture even with additional sodium acetate. In MAC-1 and MAC-2 experiments, the acetogenic bacteria inhibited also with sodium acetate influence in the medium.

The MAC-3 medium, which follows the same medium concentration of MAC-2 except sodium acetate, enhanced acetogenic bacteria and inhibited the methanogens. After 40 days of culture, the average H<sub>2</sub> volume concentration for the MAC-3 and MAC-3\* group achieved 67.2% and 59.15% as indicated by gas analysis. This amount was 33-40 times greater than that of all other experimental groups. The exogenous organic material played an important role in carbon supply.

However, organic nutrition plays an important to supply the vitamin and other microelements besides organic nutrients. Without the vitamin and microelements besides organic nutrients supply, the consortia would grow at a slower rate, such as that observed for MAC-5.

### The nutritional metabolism analysis of methanogenic consortium

Changes in MAC-4 ion concentrations were analyzed in the initial and final media. The data from this analysis revealed that the concentrations of sodium, potassium, ammonia nitrogen and magnesium were consumed microbial. In particular, 85.78% of the sodium was utilized with microbial fermentation. The medium pH changed from 6.65 to 7.32 over this period. In contrast, the concentrations of sulfate and bicarbonate increased 1.5-7 fold (Table 2). These data indicated that sodium, nitrogen, potassium and magnesium are the key elements for methanogenic consortium fermentation. However, the sulfate should be maintained at a low level in the medium.

### The cautions orthogonal analysis

From the analysis of nutrient element concentrations, it is clear that sodium, ammonia, nitrogen, potassium and magnesium play important roles in the methanogenic microbial metabolism. The orthogonal analysis experiment was designed to identify the influence of cations in bio-methane yield. The orthogonal module introduced 4 cations, (Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>) as tested with 4 different concentrations. The lowest ion concentration was set by MAC-4 and the highest reference level for cation analysis was that of the East China Sea ion concentration. The L<sub>16</sub>(4<sup>5</sup>) orthogonal table was used as an orthogonal analysis module (Table 3).

$$R_j = \text{Max}_{k_{jm}} - \text{Min}_{k_{jm}} \quad (1)$$

$$k_{jm} = \sum_{i=1}^4 y_{jmi} / 4 \quad (2)$$

Range analysis was used to indicate the affected order among factors. Range R<sub>j</sub> was calculated with module 1.

	pH	Fluoride	Chloride	Nitrate Nitrogen	Nitrite Nitrogen	Phosphate	Sulfate	Carbonate
Initial	6.65	3.70	81.58	-	-	44.67	195.80	-
final	7.32	5.18	95.90	-	-	38.60	478.40	-
	Bicarbonate							
Sodium	Ammonia Nitrogen	Potassium	Magnesium	Calcium	Water Hardness	Total Aalkalinity		
Initial	40.66	811.90	193.60	159.00	38.89	21.06	212.56	40.66
final	328.77	115.40	39.23	35.30	14.30	22.76	115.62	328.77

**Table 2:** The MAC-4 medium ion concentration changes with methanogenic consortium bio-degradation.

Serial Number	Factors (mmol/L)				Results	
	Na	Mg	Ca	K	CH <sub>4</sub>	CO <sub>2</sub>
1	23	1.3	0.5	10.7	6.65	4.71
2	23	6.7	1.7	1.7	3.92	4.27
3	23	13.0	3.4	3.4	0.92	4.03
4	23	20.0	5.1	5.1	0.46	3.23
5	130	1.3	1.7	3.4	6.59	4.84
6	130	6.7	0.5	5.1	3.85	3.56
7	130	13.0	5.1	10.7	0.99	3.31
8	130	20.0	3.4	1.7	0.38	2.94
9	260	1.3	3.4	5.1	6.49	4.47
10	260	6.7	5.1	3.4	3.76	4.73
11	260	13.0	0.5	1.7	0.57	2.47
12	260	20.0	1.7	10.7	0.34	2.84
13	390	1.3	5.1	1.7	7.01	4.86
14	390	6.7	3.4	10.7	0.30	3.90
15	390	13.0	1.7	5.1	0.47	2.28
16	390	20.0	0.5	3.4	0.30	2.67
CH <sub>4</sub>	k <sub>1j</sub>	2.99	6.69	2.84	2.97	
	k <sub>2j</sub>	2.95	2.96	2.83	2.89	
	k <sub>3j</sub>	2.79	0.74	2.02	2.82	
	k <sub>4j</sub>	2.02	0.37	3.06	2.07	
	R <sub>j</sub>	0.97	6.32	1.03	0.90	
CO <sub>2</sub>	k <sub>1j</sub>	4.06	4.72	3.35	3.64	
	k <sub>2j</sub>	3.66	4.12	3.56	4.07	
	k <sub>3j</sub>	3.63	3.02	3.84	3.39	
	k <sub>4j</sub>	3.43	2.92	4.03	3.69	
	R <sub>j</sub>	0.63	1.80	0.68	0.68	

**Table 3:** The caution orthogonal analysis table for CH<sub>4</sub> and CO<sub>2</sub>. The results represent volume percent units for both gases. The table shows the analysis of CH<sub>4</sub> and CO<sub>2</sub> ranges.

Where  $k_{jm}$  is the average result of the m level j factor (module 2), (j, m = 1, 2, 3, 4).

Where  $y_{jmi}$  is the result of the m level j factor number i data, (j, m, i = 1, 2, 3, 4).

Range analysis confirmed that for methanogenic activity, the best ion concentrations for enhancement CH<sub>4</sub> yield were Na<sub>1</sub>, Mg<sub>1</sub>, Ca<sub>4</sub> and K<sub>1</sub>, with the rank order of element effectiveness being Mg > Ca > Na > K. The most effective ion concentration for control CO<sub>2</sub> generation were Na<sub>4</sub>, Mg<sub>4</sub>, Ca<sub>1</sub>, and K<sub>3</sub>, with its ran order being Mg > Ca = K > Na.

Notable differences exist regarding the influence of CH<sub>4</sub> and CO<sub>2</sub> on yield rates as a function of cation concentrations. The CH<sub>4</sub> bio-synthesis is substantially more sensitive to element content. Maximal Mg<sup>2+</sup> concentrations to enhance methanogen metabolism processes were 1.3mmol/L, while maximal cation concentrations for Ca<sup>2+</sup>, K<sup>+</sup>, Na<sup>+</sup> were 5.1, 1.7 and 23 mmol/L, respectively.

$$F_j = \frac{4 \sum_{i=1}^4 K_{ij}^2 - (\sum_{i=1}^{16} x_i)^2}{16 \sum_{i=1}^{16} x_i^2 - (\sum_{i=1}^{16} x_i)^2} \times 5 \quad (3)$$

Variance analysis has been used to analyze the Factor Significance (F) in this system. F was calculated with module 3.

$$K_{ij} = \sum_{i=1}^4 y_{jmi} \quad (4)$$

Where  $x_i$  is the experiment results for every factor and level.

	Factor	Square of Deviance	Degree of Freedom	F Value	Significant
CH <sub>4</sub>	Na <sup>+</sup>	2.465	3	0.892	
	Mg <sup>2+</sup>	100.905	3	36.533	*
	Ca <sup>2+</sup>	2.486	3	0.900	
	K <sup>+</sup>	2.080	3	0.753	
	Error	2.762	3		
CO <sub>2</sub>	Na <sup>+</sup>	0.842	3	8.592	
	Mg <sup>2+</sup>	9.120	3	93.061	**
	Ca <sup>2+</sup>	1.079	3	11.010	*
	K <sup>+</sup>	0.954	3	9.735	
	Error	0.098	3		

**Table 4:** The variance analysis table for 4 factors with CH<sub>4</sub> and CO<sub>2</sub> data.

Based upon the F calculation, Mg<sup>2+</sup> was the significant caution factor for CH<sub>4</sub> and CO<sub>2</sub> (Table 4). This ion is particularly sensitive in inhibiting CH<sub>4</sub> metabolism processes largely for gas-coal methanogenic consortium.

## Discussion

Microbial bio-degradation of some organic compounds of coal represents a current technology available to enhance coal-bed methane. Microbial cooperation via anaerobic digestion processes results in the implementation of secondary biogenic methane generation and gas reservoirs enhancement in mining. Based on the experiments performed in this report, the following conclusions can be garnered: (1) The organic nutrient dose should be adjusted to methanogenic consortium requirements. Excessive or deficient organic nutrient support



doses adversely affecting biogas yield; (2) MAC-4 was the most effective medium in enhancing anaerobic digestion in Fushun gas-coal seam; (3) The sodium, nitrogen, potassium and magnesium are the key elements for methanogenic consortium fermentation; (4) The cation rank order of influence Fushun methanogenic consortium metabolism was  $Mg > Ca > Na > K$ , and maximal cation concentrations of  $Mg^{2+}$ ,  $Ca^{2+}$ ,  $K^+$ ,  $Na^+$  were 1.3, 5.1, 1.7 and 23 mmol/L, respectively; (5)  $Mg^{2+}$  was a particularly sensitive factor, which could inhibit methanogenic bacteria.

## Conclusion

### The nutrient concentration for gas-coal methanogenic consortium culture

The biogas generation from coal involves a complex interaction between environmental factors and biotic communities. Hydrolytic fermentative bacteria, syntrophic acetogenic bacteria, methanogenic bacteria and many other bacteria comprise the biotic community. The environment includes not only the physical factors but also the coal, coalbed water, coal-bed gas and other complex formations which we call the environment.

The medium injected into coal seam enriching nutrients of the coal seam. The series of experiments of this report confirm that nutrient media require a strict concentration of control to be effective, especially for those involving organic materials. If the organic compounds are too rich, like that modeled with our formulations of MAC-1, MAC-2 and MAC-3, the excessive levels of nutrients will adversely affect the microbial community structure. High rates growth of hydrolytic fermentative bacteria or acetogenic bacteria could inhibit methane biosynthesis. In contrast, if lack organic nutrition, such as that modeled in MAC-5, the potential for enhancing methane biosynthesis is low and the yield proceeds at a slow rate. Concentrations of nutrition required should differ as a function of: (1) the methane biosynthesis type, such as carbon dioxide reduction or acetate fermentation; (2) coal maturity grade, such as gas-coal, flame coal, bituminous coal or lignite. The exact organic nutrition required for different coal ranks needs to be identified to maximize bio-methane yield rates.

### Ion concentration for bio-methane yield enhancement

Organic biological degradation to methane is a microbial cooperation process. Fermentative bacteria initially hydrolyze complex organic compounds to acetate, longer chained fatty acids, carbon dioxide, hydrogen,  $NH_4^+$ , and  $HS^-$ . Syntrophic hydrogen-producing (proton-reducing) acetogenic bacteria reduce intermediary metabolites to acetate, carbon dioxide, and hydrogen. Hydrogen-utilizing acetogenic bacteria demethoxylate low molecular weight ligneous compounds and ferment some hydroxylated aromatic compounds. Carbon dioxide reduction methanogenic bacteria are dependent on hydrogen, produced by other bacteria, to reduce carbon dioxide or bicarbonate to methane. And acetate fermentation methanogen yields methane via acetate bio-degradation.

Different nutrients are required for different microbial activity in the community. Results from the caution orthogonal experiments revealed that cation concentrations critically influence the metabolism process of the microbial. Except for the examination of sodium, potassium, magnesium, calcium which analyzed in the cautions orthogonal analysis, the nitrogen, yeast extract, salinity and pH could have

the potential to influence the biogas synthesis. More researches to identify the effects of ions upon the coal seam biotic community are needed to reveal the microbial activity in gas-coal beds.

## Competing Interests

The authors declare that they have no competing financial interests.

## Acknowledgment

The authors acknowledge the contributions of the following companies for allowing access to coal samples and other information used in this paper: Laohutai Mining.

This work was supported by The Independent Research Project of State Key Laboratory of Coal Resources and Safe Mining, CUMT (grant number SKLCRSM19X0012: DX), The Open Research Project of State Key Laboratory of Coal Resources and Safe Mining (grant number SKLCRSM17KFA08: DX), The Fundamental Key Projects of Shanxi Provincial Key Research and Development Program (grant number 201703D211003: YZ).

## References

1. Lv Y, Tang D, Xu H, Luo H (2012) Production characteristics and the key factors in high-rank coalbed methane fields: A case study on the Fanzhuang Block, Southern Qinshui Basin, China. *Int J Coal Geol* 96: 93-108.
2. Park SY, Liang Y (2016) Biogenic methane production from coal: A review on recent research and development on microbially enhanced coalbed methane (MECBM). *Fuel*. Elsevier, UK.
3. Scott AR (1999) Improving Coal Gas Recovery with Microbially Enhanced Coalbed Methane. *Coalbed Methane: Scientific, Environmental and Economic Evaluation* 89-110.
4. Green MS, Flanagan KC, Gilcrease PC (2008) Characterization of a methanogenic consortium enriched from a coalbed methane well in the Powder River Basin USA. *Int J Coal Geol* 76: 34-45.
5. Penner TJ, Foght JM, Budwill K (2010) Microbial diversity of western Canadian subsurface coal beds and methanogenic coal enrichment cultures. *Int J Coal Geol* 82: 81-93.
6. Strapoć D, Picardal FW, Turich C, Schaperdoth I, Macalady JL, et al. (2008) Methane-producing microbial community in a coal bed of the Illinois Basin. *Appl Environ Microbiol* 74: 2424-2432.
7. Li D, Hendry P, Faiz M (2008) A survey of the microbial populations in some Australian coalbed methane reservoirs. *Int J Coal Geol* 76: 14-24.
8. McIntosh J, Martini A, Petsch S, Huang R, Nüsslein K (2008) Biogeochemistry of the Forest City Basin coalbed methane play. *Int J Coal Geol* 76: 111-118.
9. Hall RO, Tank JL, Baker MA, Rosi-Marshall EJ, Hotchkiss ER (2016) Metabolism, Gas Exchange, and Carbon Spiraling in Rivers. *Ecosystems* 19: 73-86.
10. Beckmann S, Luk AWS, Gutierrez-Zamora ML, Chong NHH, Thomas T, et al. (2018) Long-term succession in a coal seam microbiome during in situ biostimulation of coalbed-methane generation. *ISME Journal* 13: 632-650.
11. Chen T, Zheng H, Hamilton S, Rodrigues S, Golding SD, et al. (2017) Characterisation of bioavailability of Surat Basin Walloon coals for biogenic methane production using environmental microbial consortia. *Int J Coal Geol* 176: 92-112.
12. Xiao D, Wang E-Y, Peng S-P, Wu J-Y (2017) Responses of coal anaerobic fermentation fractures development. *Meitan Xuebao/Journal China Coal Soc*.

13. Meng J, Nie B, Zhao B, Ma Y (2015) Study on law of raw coal seepage during loading process at different gas pressures. *Int J Min Sci Technol* 25: 31-35.
14. Galagan JE, Nusbaum C, Roy A, Endrizzi MG, Macdonald P, et al. (2002) The genome of *M. acetivorans* reveals extensive metabolic and physiological diversity. *Genome Res* 12: 532-542.
15. Wang E, Chen P, Li Z, Shen R, Xu J, et al. (2013) Resistivity response in complete stress-strain process of loaded coal. *Meitan Xuebao/Journal China Coal Soc*.
16. Shah FA, Mahmood Q, Shah MF, Pervez A, Asad SA (2014) Microbial ecology of anaerobic digesters: The key players of anaerobiosis. *The Scientific World Journal*. 2014: 183752.
17. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, et al. (2018) A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nat Biotechnol* 36: 996-1004.
18. Zinder SH (1993) Physiological Ecology of Methanogens. *Methanogenesis* 128-206.
19. Bellou S, Baeshen MN, Elazzazy AM, Aggeli D, Sayegh F, et al (2014) Microalgal lipids biochemistry and biotechnological perspectives. *Biotechnology Advances* 32: 1476-1493.
20. Gerritsen J, Fuentes S, Grievink W, van Niftrik L, Tindall BJ, et al. (2014) Characterization of *Romboutsia ilealis* gen. nov., sp. nov., isolated from the gastro-intestinal tract of a rat, and proposal for the reclassification of five closely related members of the genus *Clostridium* into the genera *Romboutsia* gen. nov., *Intestinibacter* gen. nov., *Terrisporobacter* gen. nov. and *Asaccharospora* gen. nov. *Int J Syst Evol Microbiol* 64: 1600-1616.
21. Kwietniewska E, Tys J (2014) Process characteristics, inhibition factors and methane yields of anaerobic digestion process, with particular focus on microalgal biomass fermentation. *Renew Sustain Energy Rev* 34: 491-500.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>