

Case Report

Case Report of an Atypical Evolution of Urine Obstruction, is this Acute on Chronic Kidney Disease?

De la Vega Méndez FM¹, Alvarado Villarreal FJ¹, Purón-González E¹, Martínez Jiménez JG^{1,2,3}, and Rizo Topete L^{1,2,3*}

¹Fellow of the Internal Medicine Department, Hospital Christus Muguerza Alta Especialidad, UDEM, Monterrey, Nuevo León, México

²Internal Medicine and Nephrology Profesor at Hospital ChristusMuguerza Alta Especialidad, UDEM, Monterrey, Nuevo León, México

³Hospital Universitario "José Eleuterio González", UANL, Monterrey, Nuevo León, México

Abstract

Acute kidney injury (AKI), previously known as acute kidney failure, is defined as an abrupt (within hours) reduction of renal function in which both structural damage and loss of function are encompassed. Classification of AKI includes pre-renal, intrinsic kidney diseases (renal) and post-renal [1]. The term 'acute-on-chronic renal failure' has been used when AKI occurs in the background of pre-existing chronic kidney disease [2].

Post-renal AKI occurs after acute obstruction of the urinary flow, which increases intra-tubular pressure and thus decreases glomerular filtration rate. In addition, this acute obstruction generates an inflammatory process that contributes to the reduction of glomerular filtration. It can occur after an obstruction at any level of the urinary system (from the renal tubule to the urethra). The obstruction of the urinary tract is a common problem but a rare cause of AKI, currently there is no epidemiological data on its incidence, however it is more common in men and increases with age mainly due to prostate disease, pelvic cancer and urolithiasis [3,4]. The timing of the reversal of the obstructive cause is crucial because if not done in time can lead to permanent kidney damage. It is clear that complete or prolonged partial urinary tract obstruction can lead to tubular atrophy, interstitial fibrosis and, eventually irreversible renal injury [5]. Also is important to notice that obstruction of urinary tract may occur in the setting of preexisting Chronic Kidney Disease (CKD).

We present an atypical case (time of evolution) of a urinary obstruction with extremely high levels of plasmatic creatinine.

*Corresponding author: Rizo Topete L, Fellow of the Internal Medicine Department, Hospital Christus Muguerza Alta Especialidad, UDEM, Monterrey, Nuevo León, México, Tel: +52 1 81 8029 7718; E-mail: marili18@hotmail.com

Citation: De la Vega Méndez, Villarreal A, González P, Jiménez M, Topete R (2019) Case Report of an Atypical Evolution of Urine Obstruction, is this Acute on Chronic Kidney Disease?. J Nephrol Renal Ther 4: 022.

Received: May 13, 2019; Accepted: July 05, 2019; Published: July 12, 2019

Copyright: © 2019 De la Vega Méndez, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Keywords: Acute Kidney Injury; Postrenal Acute Kidney Injury; Urinary Tract Obstruction

Case Report

A 58-year-old man came to the emergency department with oral intolerance. He has a medical history of a perianal abscess, treated and solved. Patient refers to have presented 2 weeks before arriving to our emergency department with nausea, postprandial fullness, weight loss of eight pounds and progressive asthenia and malaise which interfered with his daily activities. The patient goes to consult where he is prescribed omeprazole for 3 weeks without improvement. Patient presented sudden vomiting and oral intolerance and arrives to our emergency department. With intentional interrogation patient refers frequency of urination, interruption of urine stream and weakness of urine stream. On physical examination: hypertensive with 177/100 mmHg, uremic breath, with presence of asterixis, oriented in time, place and person with attention preserved, without muscle weakness or sensory disturbances. Globose abdomen dull to percussion in hypogastrium, palpable mass of 15 x 13 cm in mesogastrium and hypogastrium that was mobile and not painful to palpation. Laboratories: Hemoglobin: 7.6 mg/dl, Creatinine 36.87 mg/dl, Blood urine nitrogen 211.7 mg/dl, Uric acid: 12.3 mg/dl, Albumin 4.6 g/dl Sodium: 140 mEq/L, Potassium: 6.86 mEq/L Chloride: 95.7 mEq/L. Calcium 8.6 mg/dl, Phosphorus 8.3 mg/dl. Venous Gasometry: metabolic acidosis with pH: 7.28 and HCO₃: 11.6 mEq/L, Lactate 0.5mEq/L. Urine analysis: with hemoglobin +++ and Erythrocytes 30-40 x field. An EKG was obtained in which the only relevant observed was a sinus tachycardia and QT interval prolongation (Figure 1). CT (computerized tomography) of abdomen without contrast: Hydronephrosis grade IV (Figure 2). Bladder with an estimated volume of 2 liters. Increased prostate of size 6. 5 x 5. 6 x 6 cm with an estimated volume of 115 cc that causes compression of the bladder floor.




Figure 1: Simple CT with grade IV hydronephrosis showing right kidney with measurements of 13.4 x 7.8 x 6.4 cm with increased dimensions and cortical thinning caused by the presence of hydronephrosis. Left kidney with measures of 12.3 x 7.1 x 5 cm with increased dimensions and cortical thinning caused by the presence of hydronephrosis.




Figure 2: Evolution of creatinine levels during the days of hospitalization.

The patient's obstruction was resolved on the first day of hospitalization with a urinary catheter but electrolyte abnormalities (hyperkalemia) persisted and hemodialysis was started.

Discussion

The clinical presentation of AKI due to a urinary tract obstruction varies according to the site of the obstruction, the degree of the obstruction (partial or complete) and the speed at which it occurs. It usually manifests with pain, changes in the urinary pattern, hypertension, hematuria and elevation of serum creatinine [6]. When pain is present, it is secondary to bladder distention, infection or lithiasis; hydronephrosis by itself is asymptomatic [7]. In our patient due to the large mass that was palpated in mesogastrium, it was decided to perform a CT as the first option because of the possibility of ruling out other causes of urinary retention besides increased prostate size. Although our patient had alterations in urinary volume this was not perceived by our patient as something annoying and only after an intentional interrogation is that he referred urinary symptoms. Drawing to our attention the significant elevation of the serum creatinine that normally occurs in bilateral obstruction which is the first indicator of obstruction. Obstruction of the upper and lower urinary tracts may occur in the setting of preexisting CKD whether CKD is due to an obstructive or another cause, and should be considered in all patients presenting with acute-on-chronic [8]. This may be the reason why our patient seemed to have good clinical tolerance to the considerably high levels of urea.

There are no absolute criteria for when to initiate dialysis in these patients, but it is recommended to initiate before complications occur [6,9]. It is important to remember that the serum creatinine level does not correlate with Glomerular Filtration Rate (GFR) in a patient with AKI, so the clinical manifestations will govern the decision of renal replacement therapy. It is for this reason that in the beginning this treatment was not initiated in our patient, in spite of having a potassium greater than 6.5 mEq/L and urea greater than 30mEq/L because it did not cause significant electrocardiographic changes and was not accompanied with alterations in the mental status or symptoms of severe uremic encephalopathy however despite bladder catheterization, the hyperkalemia persisted and renal replacement therapy was decided to start.

Conclusion

Although postrenal acute kidney injury is not a frequent cause of AKI could be a complication in a chronic obstruction and could drive to chronic kidney disease. The clinical presentation of an urinary obstruction should be recognized in a timely manner because it requires relatively simple treatments that can promptly restore renal function, as in this case where hydronephrosis get better but because of the length of time the diagnosis and probably the unknown chronic damage the function do not recover and the patient is still at chronic renal replacement therapy. Most of the times function recovery in AKI occurs in the first 7 to 10 days after treatment of the cause of the obstruction. However, some patients with severe renal failure may require dialysis even when the obstruction is resolved and generally creatinine levels stabilize at a level greater than 3mg/dL in those patients [9], the evolution of serum creatinine in our patient is showing in (Figure 2). Hence, it is extremely important not to forget that acute post renal kidney injury even in a setting of acute on chronic kidney disease can have severe consequences on residual renal function and urgent intervention is critical to limit renal damage and preserve residual renal function.

References

1. Makris K, Spanou L (2016) Acute Kidney Injury: Definition, Pathophysiology and Clinical Phenotypes. Clin Biochem Rev 37: 85-89.
2. Madala N (2007) Acute Renal Failure in Patients with Chronic Kidney Disease. African journalOnline 25: 395-398.
3. Bhandari S, Johnston P, Fowler RC, Joyce A, Turney JHN (1995) Non dilated bilateral ureteric obstruction. Nephrol Dial Transplant 10: 2337-2339.
4. Chapman ME, Reid JH (1991) Use of percutaneous nephrostomy in malignant ureteric obstruction. Br J Radiol 64: 318-20.
5. Blantz RC (1998) Pathophysiology of pre-renal azotemia. Kidney Int 53: 512-523.
6. Lameire N, Van Biesen W, Vanholder R (2005) Acute renal failure. The Lancet 365: 417-430.
7. Frokiaer J, Zeidel M (2011) Urinary tract obstruction. Brenner and Rector's The Kidney 9. Elsevier. New York.
8. Green J, Vardy Y, Minichor M, Better OS (1986) Extreme unilateral hydronephrosis with normal glomerular filtration rate: physiological studies in a case of obstructive uropathy. J Urol 136: 361-365.
9. Better OS, Arief AI, Massry SG, Kleeman CR, Mawell MH (1973) Studies on renal function after relief of complete unilateral ureteral obstruction of three month's duration in man. Am J Med 54: 234-240.



Journal of Anesthesia & Clinical Care
Journal of Addiction & Addictive Disorders
Advances in Microbiology Research
Advances in Industrial Biotechnology
Journal of Agronomy & Agricultural Science
Journal of AIDS Clinical Research & STDs
Journal of Alcoholism, Drug Abuse & Substance Dependence
Journal of Allergy Disorders & Therapy
Journal of Alternative, Complementary & Integrative Medicine
Journal of Alzheimer's & Neurodegenerative Diseases
Journal of Angiology & Vascular Surgery
Journal of Animal Research & Veterinary Science
Archives of Zoological Studies
Archives of Urology
Journal of Atmospheric & Earth-Sciences
Journal of Aquaculture & Fisheries
Journal of Biotech Research & Biochemistry
Journal of Brain & Neuroscience Research
Journal of Cancer Biology & Treatment
Journal of Cardiology: Study & Research
Journal of Cell Biology & Cell Metabolism
Journal of Clinical Dermatology & Therapy
Journal of Clinical Immunology & Immunotherapy
Journal of Clinical Studies & Medical Case Reports
Journal of Community Medicine & Public Health Care
Current Trends: Medical & Biological Engineering
Journal of Cytology & Tissue Biology
Journal of Dentistry: Oral Health & Cosmesis
Journal of Diabetes & Metabolic Disorders
Journal of Dairy Research & Technology
Journal of Emergency Medicine Trauma & Surgical Care
Journal of Environmental Science: Current Research
Journal of Food Science & Nutrition
Journal of Forensic, Legal & Investigative Sciences
Journal of Gastroenterology & Hepatology Research
Journal of Gerontology & Geriatric Medicine
Journal of Genetics & Genomic Sciences
Journal of Hematology, Blood Transfusion & Disorders
Journal of Human Endocrinology
Journal of Hospice & Palliative Medical Care
Journal of Internal Medicine & Primary Healthcare
Journal of Infectious & Non Infectious Diseases
Journal of Light & Laser: Current Trends
Journal of Modern Chemical Sciences
Journal of Medicine: Study & Research
Journal of Nanotechnology: Nanomedicine & Nanobiotechnology
Journal of Neonatology & Clinical Pediatrics
Journal of Nephrology & Renal Therapy
Journal of Non Invasive Vascular Investigation
Journal of Nuclear Medicine, Radiology & Radiation Therapy
Journal of Obesity & Weight Loss
Journal of Orthopedic Research & Physiotherapy
Journal of Otolaryngology, Head & Neck Surgery
Journal of Protein Research & Bioinformatics
Journal of Pathology Clinical & Medical Research
Journal of Pharmacology, Pharmaceutics & Pharmacovigilance
Journal of Physical Medicine, Rehabilitation & Disabilities
Journal of Plant Science: Current Research
Journal of Psychiatry, Depression & Anxiety
Journal of Pulmonary Medicine & Respiratory Research
Journal of Practical & Professional Nursing
Journal of Reproductive Medicine, Gynaecology & Obstetrics
Journal of Stem Cells Research, Development & Therapy
Journal of Surgery: Current Trends & Innovations
Journal of Toxicology: Current Research
Journal of Translational Science and Research
Trends in Anatomy & Physiology
Journal of Vaccines Research & Vaccination
Journal of Virology & Antivirals
Archives of Surgery and Surgical Education
Sports Medicine and Injury Care Journal
International Journal of Case Reports and Therapeutic Studies

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>