



Mini Review

## Role of Rehabilitation in Innovative Models for Mobility in the Short-Term Care Setting

Rebecca J Tarbert\*

Director of Clinical Programs, Active Protective Technologies, Inc, Fort Washington, Pennsylvania, USA

The role of rehabilitation services in the short-term care unit is shifting along with the inception of value-based care models in healthcare. Healthcare organizations seeking efficiency and effectiveness for reduced spending and decreased risk of re-hospitalization are looking to achieve meaningful outcomes using evidence-based care models through interdisciplinary approaches [1]. In the United States specifically, the expectation of the Patient Driven Payment Model is an example of reimbursement for health care services delivered received based upon accuracy of patient need assessment and change in functional status. The amount of time a patient spends with their physical, occupational and speech language pathologist each day is no longer the driving equation of how much the facility is reimbursed for the care provided. In this new value based care model, the care needs specific to the patient's condition are weighted to provide reimbursement that is based upon the outcome of the care provided [2]. Rehabilitation services will continue to be an integral piece of achieving success in the PDPM landscape with continued focus on remediation, compensation and education techniques our therapists have always provided. What the shift warrants for rehabilitation services is the increased need for interprofessional collaboration and focused efficiency to provide better care and improved outcomes at a lower cost in the post-acute setting of short-term rehabilitation.

Rehabilitation services provide a critical aspect of care in short-term rehab in screening, evaluating and providing rehabilitation intervention to patients admitted in their facilities to return to home. A key part of the rehabilitation stay is the patient's participation in mobility while in the center, balanced with any fall risk factors that may be present. Rehabilitation holds a role of identifying the physical and cognitive ability of patients to participate in mobility performance while in the short stay setting [3]. Gait performance, activity

\*Corresponding author: Rebecca J Tarbert, Director of Clinical Programs, Active Protective Technologies, Inc, Fort Washington, Pennsylvania, USA, Tel: +1 2672426125; E-mail: rebecca@activeprotective.com

Citation: Tarbert RJ (2019) Role of Rehabilitation in Innovative Models for Mobility in the Short-Term Care Setting. J Phys Med Rehabil Disabil 5: 035.

Received: August 01, 2019; Accepted: September 17, 2019; Published: September 24, 2019

Copyright: © 2019 Tarbert RJ. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

tolerance, strength, balance, and safety awareness are all components of the assessment as to whether a patient is able to perform a level of mobility without assistance while staying in the care facility [4].

While the implementation of fall mitigation strategies and equipment for those patients identified as being at high risk of falling are intentioned to provide maximum safety to the patient, many times the strategies lead to limitations in the patient's ability to mobilize independently. It is well established that increased participation in physical activity by older adults in the community as well as while in in-patient care settings not only maintains functional status, but advances the ability to perform functional activities and reduces risk of adverse events such as falls and hospital re-admissions [5-9]. The challenge in the multifactorial setting of short-term rehab is balancing how to empower patients to participate in mobility while maintaining safety during their physically vulnerable state. Barriers to increasing mobility outside of scheduled therapy time frames stem from not only rehabilitation department burdens, but the many tasks all members of the interdisciplinary team perform. Most research on the barriers associated with rehabilitation and nursing care to facilitate increased patient mobility are rooted in hospital care, though findings can be related to the challenging in-patient setting of short-term care. Barriers noted include having enough time to offer mobilization, knowledge of how to provide mobility safely to patients, low physical resources, low staff resources and therapist caseloads for skilled intervention versus offering repetitive practiced mobility activities [10,11].

In the skilled nursing environment, the utilization of restorative care to offer mobility participation for in-patient residents has been utilized primarily for the long-term care resident maintain ADL function and quality of life [12]. In the incoming PDPM model of care, restorative care is able to be captured for long-term as well as short-term care needs of patients as the role of physical activity outside of usual rehabilitation intervention is recognized to benefit the patient experience and support patient functional outcomes. A literature review by Benjamin et al. in 2014 revealed that the implementation of mobility programming in the skilled nursing environment required an interdisciplinary approach addressing the funding and staffing constraints shared by most of the randomized controlled trials [13]. A call to action was issued for care providers to consider innovative models of mobility in the in-patient setting recognizing socioeconomic and organizational barriers to facilitate mobility in every day care.

Innovative models of interdisciplinary collaboration focused on patient-specific goals are being explored in the skilled nursing industry as organizations seek to find efficient and effective ways to achieve value-based care. Creative partnerships of care providers in and between organizations consider how care is delivered and overseen in a way that most benefits the patient. Many organizations are evaluating upcoming technologies that can increase participation of patient mobility while decreasing burden on care teams, thereby allowing the focused care patients need. Technology is allowing for improved communication between providers, increased access to care in all settings, assisted interventions and identification of evidence-

based protocols in care delivery [14]. In the goal of addressing safety in the short-term care setting, technology is utilized in documentation systems, communication between providers and in fall risk management. Traditional technologies for fall risk management and injury prevention in short-stay settings include the use of bed/chair sensors to alert nursing staff when a patient is moving out of the stationary position and personal emergency alert systems that notify nursing staff when a patient needs assistance. These technologies support communication to caregivers but have not been proven to decrease falls or injuries from falls and have been found to increase burden to care teams and limit patient participation in mobility [15].

Emerging technologies in support of the safe performance of mobility include the implementation of a smart hip protection belt. The light, form-fitting belt can always be worn when a patient is in the short stay center to offer protection to the proximal femur using fall detection and intervention technology. In the event of a serious hip-impacting fall, the belt will deploy airbags around the hips to attenuate impact forces and alert caregivers as to the need for assistance [16].

Fall risk factors that are associated with a high risk of hip fracture include the level of mobility of patients who are receiving care in short-term rehab. These patients are attending to rehabilitation as they are not quite strong enough to perform activities of daily living completely independently or safely, though they are strengthening and returning to an independent level. Those patients who are able to perform a transfer, but demonstrate limited ability to maintain balance and/or perform locomotion once standing, are those at highest risk of hip fracture from falls [17,18]. Rehabilitation professionals assessing the mobility level with each patient in the setting and continuing to re-assess with each daily visit can identify those who may benefit from wearable hip protection and offer decreased risk of fall injury while in this transitional phase.

Implementation of this new technology has demonstrated an increased participation in mobility of the short-term care patients who did wear the device by 50% as compared to those who did not choose to wear it. To explore underlying reasons for increased mobility associated with use of the hip protection device, a concern-of-falling survey was administered to belt wearers and found that those who wore the hip protection belt while undergoing short-term rehabilitation reported a 22% increase in their confidence surrounding the performance of daily movements and activities. This early finding is consistent with others who have shown a correlation between fear of falling and self-limiting of mobility [19,20]. The relationship between a decrease in concern-of-falling and the increased participation in mobility while utilizing a smart hip protection belt during a rehabilitation stay merits further investigation.

Rehabilitation professionals can champion a culture of mobility with the use of technology which offers safe mobility to the patient in the short-term care setting. This highlights their role of prescribing mobility, maintaining safety and supporting patient-driven goals for reducing risk of falling, increasing functional outcomes and supporting discharge planning. Advancements of the wearable technology to be evaluated in the future in support of patient and rehabilitation goals are: mobility in step count per day, postural sway (deviation from vertical axis) in the performance of daily activities, on demand testing of Timed up and Go, gait speed and repeated Short FES-I surveys.

Access to the mobility metrics by the interdisciplinary team via mobile application or desktop will provide interdisciplinary communication and foster collaboration for patient-centered decision making. The utilization of such measures with easy access to tracking and data sharing further enhances the quality reporting for outcomes of rehabilitative care for the short-term facility [21].

Interdisciplinary collaboration with patient driven goals focused on their specific needs to function independently and return to home are critical in rehabilitation settings. With each care discipline recognizing their role in the collaborative care setting and also recognizing the importance of clear communication on shared goals, the utilization of technology has the capability of reducing burdens on staff for enabling safe mobility while improved functional outcomes in the short-stay setting [22,23]. Rehabilitation providers have the opportunity now as they hold a unique skill set, to champion innovative programs and technologies to achieve value-based outcomes compelling their role in post-acute care.

## References

1. Reyes B, Ouslander JG (2019) *Eur Geriatr Med* 10: 387.
2. Centers for Medicare and Medicaid Services 2019, Patient Driven Payment Model. Accessed 7/16/2019.
3. Muir SW, Gopaul K, Montero Odasso MM (2012) The role of cognitive impairment in fall risk among older adults: a systematic review and meta-analysis. *Age and Ageing* 41: 299-308.
4. Karen Hertz, Julie Santy Tomlinson (2018) Fragility fracture nursing, perspectives in nursing management and care for older adults. Pg no: 155-158.
5. Zisberg A, Shadmi E, Sinoff G, Gur-Yaish N, Sruлович E, et al. (2011) Low mobility during hospitalization and functional decline in older adults. *J Am Geriatr Soc* 59: 266-273.
6. Merom D, Pye V, Macniven R, van der Ploeg H, Milat A, et al. (2012) Prevalence and correlates of participation in fall prevention exercise/physical activity by older adults. *Prev Med* 55: 613-617.
7. Sherrington C, Tiedmann A, Cameron I (2011) Physical exercise after hip fracture: An evidence overview. *Eur J Phys Rehabil Med* 47: 297-307.
8. Fisher SR, Kuo YF, Sharma G, Raji MA, Kumar A, et al. (2013) Mobility after hospital discharge as a marker for 30-day readmission. *J Gerontol A BiolSci Med Sci* 68: 805-810.
9. Hoyer EH, Friedman M, Lavezza A, Wagner-Kosmakos K, Lewis-Cherry R, et al. (2016) Promoting mobility and reducing length of stay in hospitalized general medicine patients: A quality-improvement project. *J Hosp Med* 11: 341-347.
10. Hoyer EH, Brotman DJ, Chan KS, Needham DM (2015) Barriers to early mobility of hospitalized general medicine patients: survey development and results. *Am J Phys Med Rehabil* 94: 304-312.
11. Coughenour ET, Salmans KJ, Skoch AD, Starks LL, Sabus, C (2017) Achieving a culture of mobility: Implementation of a mobility aide program to increase patient mobilizations in an acute care hospital. *J Acute Care Phys Ther* 8: 86-95.
12. Remsburg RE, Armacost KA, Radu C, Bennett RG (2001) Impact of a restorative care program in the nursing home. *Educ Gerontol* 27: 261-280.
13. Benjamin K, Edwards N, Ploeg J, Legault F (2014) Barriers to physical activity and restorative care for residents in long-term care: a review of the literature. *J Aging Phys Act* 22: 154-165.

14. Dicianno BE, Parmanto B, Fairman AD, Crytzer TM, Yu DX, et al. (2015) Perspectives on the evolution of mobile (mHealth) technologies and application to rehabilitation. *Phys Ther* 95:397-405.
15. Shorr RI, Chandler AM, Mion LC, Waters TM, Liu M, et al. (2012) Effects of an intervention to increase bed alarm use to prevent falls in hospitalized patients: a cluster randomized trial. *Ann Intern Med* 157: 692-699.
16. Quigley P, Singhatat W, Tarbert R (2019) Technology innovation to protect hips from fall-related fracture. *Phys Med Rehabil Res* 4: 1-4.
17. Chen JS, Sambrook PN, Simpson JM, Cameron ID, Cumming RG, et al. (2009) Risk factors for hip fracture among institutionalised older people. *Age and Ageing* 38: 429-434.
18. Jaglal SB, Hawker G, Cameron C, Canavan J, Beaton D, et al. (2010) The ontario osteoporosis strategy: implementation of a population-based osteoporosis action plan in canada. *Osteoporos Int* 21: 903-908.
19. Friedman SM, Munoz B, West SK, Rubin GS, Fried LP (2002) Falls and fear of falling: which comes first? a longitudinal prediction model suggests strategies for primary and secondary prevention. *J Am Geriatr Soc* 50: 1329-1335.
20. Gaxatte C, Nguyen T, Chourabi F, Salleron J, Pardessus V, et al. (2011) Fear of falling as seen in the multidisciplinary falls consultation. *Ann Phys Rehabil Med* 54: 248-258.
21. Horak F, King L, Mancini M (2015) Role of body-worn movement monitor technology for balance and gait rehabilitation. *Phys Ther* 95: 461-470.
22. May CR (2015) Making sense of technology adoption in healthcare: meso-level considerations. *BMC Medicine* 13 (92).
23. Jensen RE, Rothrock NE, DeWitt EM, Spiegel B, Tucker CA, et al. (2015) The role of technical advances in the adoption and integration of patient-reported outcomes in clinical care. *Med Care* 53:153-159.



Journal of Anesthesia & Clinical Care  
Journal of Addiction & Addictive Disorders  
Advances in Microbiology Research  
Advances in Industrial Biotechnology  
Journal of Agronomy & Agricultural Science  
Journal of AIDS Clinical Research & STDs  
Journal of Alcoholism, Drug Abuse & Substance Dependence  
Journal of Allergy Disorders & Therapy  
Journal of Alternative, Complementary & Integrative Medicine  
Journal of Alzheimer's & Neurodegenerative Diseases  
Journal of Angiology & Vascular Surgery  
Journal of Animal Research & Veterinary Science  
Archives of Zoological Studies  
Archives of Urology  
Journal of Atmospheric & Earth-Sciences  
Journal of Aquaculture & Fisheries  
Journal of Biotech Research & Biochemistry  
Journal of Brain & Neuroscience Research  
Journal of Cancer Biology & Treatment  
Journal of Cardiology: Study & Research  
Journal of Cell Biology & Cell Metabolism  
Journal of Clinical Dermatology & Therapy  
Journal of Clinical Immunology & Immunotherapy  
Journal of Clinical Studies & Medical Case Reports  
Journal of Community Medicine & Public Health Care  
Current Trends: Medical & Biological Engineering  
Journal of Cytology & Tissue Biology  
Journal of Dentistry: Oral Health & Cosmesis  
Journal of Diabetes & Metabolic Disorders  
Journal of Dairy Research & Technology  
Journal of Emergency Medicine Trauma & Surgical Care  
Journal of Environmental Science: Current Research  
Journal of Food Science & Nutrition  
Journal of Forensic, Legal & Investigative Sciences  
Journal of Gastroenterology & Hepatology Research  
Journal of Gerontology & Geriatric Medicine  
Journal of Genetics & Genomic Sciences  
Journal of Hematology, Blood Transfusion & Disorders  
Journal of Human Endocrinology  
Journal of Hospice & Palliative Medical Care  
Journal of Internal Medicine & Primary Healthcare  
Journal of Infectious & Non Infectious Diseases  
Journal of Light & Laser: Current Trends  
Journal of Modern Chemical Sciences  
Journal of Medicine: Study & Research  
Journal of Nanotechnology: Nanomedicine & Nanobiotechnology  
Journal of Neonatology & Clinical Pediatrics  
Journal of Nephrology & Renal Therapy  
Journal of Non Invasive Vascular Investigation  
Journal of Nuclear Medicine, Radiology & Radiation Therapy  
Journal of Obesity & Weight Loss  
Journal of Orthopedic Research & Physiotherapy  
Journal of Otolaryngology, Head & Neck Surgery  
Journal of Protein Research & Bioinformatics  
Journal of Pathology Clinical & Medical Research  
Journal of Pharmacology, Pharmaceutics & Pharmacovigilance  
Journal of Physical Medicine, Rehabilitation & Disabilities  
Journal of Plant Science: Current Research  
Journal of Psychiatry, Depression & Anxiety  
Journal of Pulmonary Medicine & Respiratory Research  
Journal of Practical & Professional Nursing  
Journal of Reproductive Medicine, Gynaecology & Obstetrics  
Journal of Stem Cells Research, Development & Therapy  
Journal of Surgery: Current Trends & Innovations  
Journal of Toxicology: Current Research  
Journal of Translational Science and Research  
Trends in Anatomy & Physiology  
Journal of Vaccines Research & Vaccination  
Journal of Virology & Antivirals  
Archives of Surgery and Surgical Education  
Sports Medicine and Injury Care Journal  
International Journal of Case Reports and Therapeutic Studies  
Journal of Ecology Research and Conservation Biology

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>